

CS-184: Computer Graphics

Lecture #12: Curves and Surfaces

Prof. James O'Brien
University of California, Berkeley

V2009-F-12.1.0

Today


- General curve and surface representations
- Splines and other polynomial bases

Geometry Representations

- Constructive Solid Geometry (CSG)
- Parametric
 - Polygons
 - Subdivision surfaces
- Implicit Surfaces
- Point-based Surface
- Not always clear distinctions
 - *i.e.* **CSG done with implicits**

3

Geometry Representations




Object made by CSG
Converted to
polygons

4

Geometry Representations


Object made by CSG
Converted to polygons
Converted to implicit surface



5


Geometry Representations

CSG on implicit surfaces




6

Geometry Representations







Point-based surface descriptions



Ohtake, *et al.*, SIGGRAPH 2003

Geometry Representations



Subdivision surface
(different levels of refinement)

Images from Subdivision.org

Geometry Representations

- Various strengths and weaknesses
 - Ease of use for design
 - Ease/speed for rendering
 - Simplicity
 - Smoothness
 - Collision detection
 - Flexibility (in more than one sense)
 - Suitability for simulation
 - *many others...*

9

Parametric Representations

Curves: $\mathbf{x} = \mathbf{x}(u)$ $\mathbf{x} \in \mathbb{R}^n$ $u \in \mathbb{R}$

Surfaces: $\mathbf{x} = \mathbf{x}(u, v)$ $\mathbf{x} \in \mathbb{R}^n$ $u, v \in \mathbb{R}$
 $\mathbf{x} = \mathbf{x}(\mathbf{u})$ $\mathbf{u} \in \mathbb{R}^2$

Volumes: $\mathbf{x} = \mathbf{x}(u, v, w)$ $\mathbf{x} \in \mathbb{R}^n$ $u, v, w \in \mathbb{R}$
 $\mathbf{x} = \mathbf{x}(\mathbf{u})$ $\mathbf{u} \in \mathbb{R}^3$


and so on...

Note: a vector function is really n scalar functions


10

Parametric Rep. Non-unique

- Same curve/surface may have multiple formulae



$$\mathbf{x}(u) = [u, u]$$



$$\mathbf{x}(u) = [u^3, u^3]$$

11

Simple Differential Geometry

- Tangent to curve

$$\mathbf{t}(u) = \frac{\partial \mathbf{x}}{\partial u} \Big|_u$$

- Tangents to surface


$$\mathbf{t}_u(u, v) = \frac{\partial \mathbf{x}}{\partial u} \Big|_{u,v} \quad \mathbf{t}_v(u, v) = \frac{\partial \mathbf{x}}{\partial v} \Big|_{u,v}$$

- Normal of surface

$$\hat{\mathbf{n}} = \frac{\mathbf{t}_u \times \mathbf{t}_v}{\|\mathbf{t}_u \times \mathbf{t}_v\|}$$

- Also: curvature, curve normals, curve bi-normal, **others...**


- Degeneracies: $\partial \mathbf{x} / \partial u = 0$ or $\mathbf{t}_u \times \mathbf{t}_v = 0$



12

Discretization

- Arbitrary curves have an uncountable number of parameters



i.e. specify function value at all points on real number line

13

Discretization

- Arbitrary curves have an uncountable number of parameters

- Pick **complete** set of basis functions

$$x(u) = \sum_{i=0}^{\infty} c_i \phi_i(u)$$

- Polynomials, Fourier series, **etc.**

- Truncate set at some reasonable point

$$x(u) = \sum_{i=0}^3 c_i \phi_i(u) = \sum_{i=0}^3 c_i u^i$$

- Function represented by the vector (list) of c_i

- The c_i may themselves be vectors

$$\mathbf{x}(u) = \sum_{i=0}^3 \mathbf{c}_i \phi_i(u)$$

14

Polynomial Basis

- Power Basis

$$x(u) = \sum_{i=0}^d c_i u^i$$

$$x(u) = \mathbf{C} \cdot \mathcal{P}^d$$

$$\mathbf{C} = [c_0, c_1, c_2, \dots, c_d]$$

$$\mathcal{P}^d = [1, u, u^2, \dots, u^d]$$

The elements of \mathcal{P}^d are **linearly independent**

i.e. no good approximation

$$u^k \neq \sum_{i \neq k} c_i u^i$$


Skipping something would lead to bad results... odd stiffness

15

Specifying a Curve

Given desired values (constraints) how do we determine the coefficients for cubic power basis?

For now, assume
 $u_0 = 0 \quad u_1 = 1$



16

Specifying a Curve


Given desired values (constraints) how do we determine the coefficients for cubic power basis?

$$x(0) = c_0 = x_0$$

$$x(1) = \sum c_i = x_1$$

$$x'(0) = c_1 = x'_0$$

$$x'(1) = \sum i c_i = x'_1$$




17

Specifying a Curve

Given desired values (constraints) how do we determine the coefficients for cubic power basis?

$$\begin{bmatrix} x_0 \\ x_1 \\ x'_0 \\ x'_1 \end{bmatrix} = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 1 & 1 & 1 & 1 \\ 0 & 1 & 0 & 0 \\ 0 & 1 & 2 & 3 \end{bmatrix} \cdot \begin{bmatrix} c_0 \\ c_1 \\ c_2 \\ c_3 \end{bmatrix}$$

$$\mathbf{p} = \mathbf{B} \cdot \mathbf{c}$$




18

Specifying a Curve

Given desired values (constraints) how do we determine the coefficients for cubic power basis?

$$\mathbf{c} = \beta_H \cdot \mathbf{p}$$

$$\beta_H = \mathbf{B}^{-1} = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ -3 & 3 & -2 & 1 \\ 2 & -2 & 1 & 1 \end{bmatrix}$$



19


Specifying a Curve

Given desired values (constraints) how do we determine the coefficients for cubic power basis?

$$\mathbf{c} = \beta_H \cdot \mathbf{p}$$

$$x(u) = \mathcal{P}^3 \cdot \mathbf{c} = \boxed{\mathcal{P}^3 \beta_H} \mathbf{p}$$

$$= \begin{bmatrix} 1 + 0u - 3u^2 + 2u^3 \\ 0 + 0u + 3u^2 - 2u^3 \\ 0 + 1u - 2u^2 + 1u^3 \\ 0 + 0u - 1u^2 + 1u^3 \end{bmatrix} \mathbf{p}$$



$$\beta_H = \mathbf{B}^{-1} = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ -3 & 3 & -2 & 1 \\ 2 & -2 & 1 & 1 \end{bmatrix}$$


20

Specifying a Curve

Given desired values (constraints) how do we determine the coefficients for cubic power basis?

$$\mathbf{c} = \beta_H \cdot \mathbf{p}$$

$$x(u) = \begin{bmatrix} 1 + 0u - 3u^2 + 2u^3 \\ 0 + 0u + 3u^2 - 2u^3 \\ 0 + 1u - 2u^2 + 1u^3 \\ 0 + 0u - 1u^2 + 1u^3 \end{bmatrix} \mathbf{p}$$





$$x(u) = \sum_{i=0}^3 p_i b_i(u)$$

Hermite basis functions

Specifying a Curve

Given desired values (constraints) how do we determine the coefficients for cubic power basis?



$$x(u) = \sum_{i=0}^3 p_i b_i(u)$$

Hermite basis functions

Hermite Basis


- Specify curve by
 - Endpoint values
 - Endpoint tangents (derivatives)
- Parameter interval is arbitrary (most times)
 - Don't need to recompute basis functions
- These are **cubic** Hermite
 - Could do construction for any odd degree
 - $(d - 1)/2$ derivatives at end points

23

Cubic Bézier

- Similar to Hermite, but specify tangents indirectly

$$\begin{aligned}x_0 &= p_0 \\x_1 &= p_3 \\x'_0 &= 3(p_1 - p_0) \\x'_1 &= 3(p_3 - p_2)\end{aligned}$$



Note: all the control points are points in space, no tangents.

24


Cubic Bézier

- Similar to Hermite, but specify tangents indirectly

$$\begin{bmatrix} 1 & 0 & 0 & 0 \\ 1 & 1 & 1 & 1 \\ 0 & 1 & 0 & 0 \\ 0 & 1 & 2 & 3 \end{bmatrix} \mathbf{c} = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 \\ -3 & 3 & 0 & 0 \\ 0 & 0 & -3 & 3 \end{bmatrix} \mathbf{p}$$

$$\mathbf{c} = \begin{bmatrix} 1 & 0 & 0 & 0 \\ -3 & 3 & 0 & 0 \\ 3 & -6 & 3 & 0 \\ -1 & 3 & -3 & 1 \end{bmatrix} \mathbf{p}$$


$$\mathbf{c} = \beta_z \mathbf{p}$$



25

Cubic Bézier

- Plot of Bézier basis functions



26

Changing Bases

- Power basis, Hermite, and Bézier all are still just cubic polynomials

- The three basis sets all span the same space
- Like different axes in

- Changing basis $\mathbb{R}^3 \times \mathbb{R}^4$

$$\mathbf{c} = \beta_Z \mathbf{p}_Z$$

$$\mathbf{c} = \beta_H \mathbf{p}_H$$

$$\mathbf{p}_Z = \beta_Z^{-1} \beta_H \mathbf{p}_H$$


27

Useful Properties of a Basis

- Convex Hull

- All points on curve inside convex hull of control points
- Bézier basis has convex hull property

$$\sum_i b_i(u) = 1 \quad b_i(u) \geq 0 \quad \forall u \in \Omega$$



28

Useful Properties of a Basis

- Invariance under class of transforms
 - Transforming curve is same as transforming control points
 - Bézier basis invariant for affine transforms
 - Bézier basis NOT invariant for perspective transforms
 - NURBS are though...

$$\mathbf{x}(u) = \sum_i \mathbf{p}_i b_i(u) \Leftrightarrow \mathcal{T}\mathbf{x}(u) = \sum_i (\mathcal{T}\mathbf{p}_i) b_i(u)$$

29


Useful Properties of a Basis

- Local support
 - Changing one control point has limited impact on entire curve
- Nice subdivision rules
- Orthogonality ($\int_{\Omega} b_i(u) b_j(u) du = \delta_{ij}$)
- Fast evaluation scheme
- Interpolation -vs- approximation


30

DeCasteljau Evaluation

- A geometric evaluation scheme for Bézier



DeCasteljau Evaluation




From Wikipedia

Adaptive Tessellation

- Midpoint test subdivision
- Possible problem
 - Simple solution if curve basis has **convex hull** property



If curve inside convex hull and the convex hull is nearly flat: curve is nearly flat and can be drawn as straight line




Better: draw convex hull

Works for Bézier because the ends are interpolated

33

Bézier Subdivision


- Form control polygon for half of curve by evaluating at $u=0.5$



34

Bézier Subdivision

- Form control polygon for half of curve by evaluating at $u=0.5$



34


Bézier Subdivision

- Form control polygon for half of curve by evaluating at $u=0.5$

Repeated subdivision
makes smaller/flatter
segments


Also works for surfaces...

We'll extend this idea
later on...



34

Joining



$$c^0 \Leftrightarrow b = b$$

$$c^1 \Leftrightarrow b - a = c - b$$

$$G^1 \Leftrightarrow \frac{b - a}{\|b - a\|} = \frac{c - b}{\|c - b\|}$$


If you change **a**, **b**, or **c** you must change the others

But if you change **a**, **b**, or **c** you do not have to change beyond those three. *LOCAL SUPPORT*

35

“Hump” Functions

- Constraints at joining can be built in to make new basis



36

Tensor-Product Surfaces

- Surface is a curve swept through space
- Replace control points of curve with other curves


$$x(u, v) = \sum_i p_i b_i(u) \quad q_i(v) = \sum_j p_{ji} b_j(v)$$

$$x(u, v) = \sum_{ij} p_{ij} b_i(u) b_j(v) \quad b_{ij}(u, v) = b_i(u) b_j(v)$$

$$x(u, v) = \sum_{ij} p_{ij} b_{ij}(u, v)$$


37

Tensor-Product Surfaces




38


Hermite Surface Bases



Hermite Surface Hump




Bézier Surface Patch




Adaptive Tessellation

- Given surface patch
 - If close to flat: draw it
 - Else subdivide 4 ways



Adaptive Tessellation


- Avoid cracking



43

Adaptive Tessellation

- Avoid cracking




Cracks may be okay in some contexts...

44

Adaptive Tessellation


- Avoid cracking



45

Adaptive Tessellation

- Avoid cracking




Test interior and boundary of patch
Split boundary based on boundary
test
Table of polygon patterns
May wish to avoid "slivers"

46

Adaptive Tessellation


- Triangle Based Method (no cracks)



47

Adaptive Tessellation


- Triangle Based Method (no cracks)



48


Adaptive Tesselation

- Triangle Based Method (no cracks)



Adaptive Tesselation


- Triangle Based Method (no cracks)



$$\|B((u_1 + u_2)/2) - (x_1 + x_2)/2\| < \tau ?$$


Adaptive Tessellation

- Triangle Based Method (no cracks)



Center test tends to generate slivers.
Often better to leave it out.

Adaptive Tessellation



Without center test



With center test

Adaptive Tessellation




Second row shows typical error of swapping tests.

Yiding Jia, CS184 508 -- I broke his code to make this example.

53

Adaptive Tessellation



Visible artifacts from cracks.

Apollo Ellis, CS184 508

54