


# CS70: Jean Walrand: Lecture 24.

## Markov Chains


1. Examples
2. Definition
3. First Passage Time

## Two-State Markov Chain

Here is a symmetric two-state Markov chain. It describes a random motion in  $\{0, 1\}$ . Here,  $a$  is the probability that the state changes in the next step.




Let's simulate the Markov chain:




# Five-State Markov Chain


At each step, the MC follows one of the outgoing arrows of the current state, with equal probabilities.



Let's simulate the Markov chain:



# Finite Markov Chain: Definition



- ▶ A finite set of states:  $\mathcal{X} = \{1, 2, \dots, K\}$
- ▶ A probability distribution  $\pi_0$  on  $\mathcal{X}$  :  $\pi_0(i) \geq 0, \sum_i \pi_0(i) = 1$
- ▶ Transition probabilities:  $P(i, j)$  for  $i, j \in \mathcal{X}$

$$P(i, j) \geq 0, \forall i, j; \sum_j P(i, j) = 1, \forall i$$

- ▶  $\{X_n, n \geq 0\}$  is defined so that

$$Pr[X_0 = i] = \pi_0(i), i \in \mathcal{X} \text{ (initial distribution)}$$


$$Pr[X_{n+1} = j | X_0, \dots, X_n = i] = P(i, j), i, j \in \mathcal{X}.$$

# First Passage Time - Example 1

Let's flip a coin with  $Pr[H] = p$  until we get  $H$ . How many flips, on average?


Let's define a Markov chain:

- ▶  $X_0 = S$  (start)
- ▶  $X_n = S$  for  $n \geq 1$ , if last flip was  $T$  and no  $H$  yet
- ▶  $X_n = E$  for  $n \geq 1$ , if we already got  $H$  (end)



## First Passage Time - Example 1

Let's flip a coin with  $Pr[H] = p$  until we get  $H$ . How many flips, on average?



Let  $\beta(S)$  be the average time until  $E$ , starting from  $S$ .

Then,

$$\beta(S) = 1 + q\beta(S) + p \cdot 0.$$


(See next slide.) Hence,

$$p\beta(S) = 1, \text{ so that } \beta(S) = 1/p.$$

Note: Time until  $E$  is  $G(p)$ . We have rediscovered that the mean of  $G(p)$  is  $1/p$ .

## First Passage Time - Example 1

Let's flip a coin with  $Pr[H] = p$  until we get  $H$ . How many flips, on average?



Let  $\beta(S)$  be the average time until  $E$ .

Then,

$$\beta(S) = 1 + q\beta(S) + p0.$$

**Justification:** Let  $N$  be the random number of steps until  $E$ , starting from  $S$ . Let also  $N'$  be the number of steps until  $E$ , after the second visit to  $S$ . Finally, let  $Z = 1\{\text{first flip} = H\}$ . Then,

$$N = 1 + (1 - Z) \times N' + Z \times 0.$$

Now,  $Z$  and  $N'$  are independent. Also,  $E[N'] = E[N] = \beta(S)$ . Hence, taking expectation,

$$\beta(S) = E[N] = 1 + (1 - p)E[N'] + p0 = 1 + q\beta(S) + p0.$$

## First Passage Time - Example 2

Let's flip a coin with  $Pr[H] = p$  until we get two consecutive  $H$ s. How many flips, on average?

*H T H T T T H T H T H T T H T H H*


Let's define a Markov chain:

- ▶  $X_0 = S$  (start)
- ▶  $X_n = E$ , if we already got two consecutive  $H$ s (end)
- ▶  $X_n = T$ , if last flip was  $T$  and we are not done
- ▶  $X_n = H$ , if last flip was  $H$  and we are not done



## First Passage Time - Example 2

Let's flip a coin with  $Pr[H] = p$  until we get two consecutive  $H$ s. How many flips, on average? Here is a picture:



$S$ : Start

$H$ : Last flip =  $H$

$T$ : Last flip =  $T$

$E$ : Done

Let  $\beta(i)$  be the average time from state  $i$  until the MC hits state  $E$ .

We claim that (these are called the [first step equations](#))


$$\beta(S) = 1 + p\beta(H) + q\beta(T)$$

$$\beta(H) = 1 + p\beta(H) + q\beta(T)$$

$$\beta(T) = 1 + p\beta(H) + q\beta(T).$$

Solving, we find  $\beta(S) = 2 + 3qp^{-1} + q^2p^{-2}$ . (E.g.,  $\beta(S) = 6$  if  $p = 1/2$ .)

## First Passage Time - Example 2



*S*: Start

*H*: Last flip = *H*

*T*: Last flip = *T*

*E*: Done

Let us justify the first step equation for  $\beta(T)$ . The others are similar.

Let  $N(T)$  be the random number of steps, starting from  $T$  until the MC hits  $E$ . Let also  $N(H)$  be defined similarly. Finally, let  $N'(T)$  be the number of steps after the second visit to  $T$  until the MC hits  $E$ . Then,

$$N(T) = 1 + Z \times N(H) + (1 - Z) \times N'(T)$$

where  $Z = 1_{\{\text{first flip in } T \text{ is } H\}}$ . Since  $Z$  and  $N(H)$  are independent, and  $Z$  and  $N'(T)$  are independent, taking expectations, we get


$$E[N(T)] = 1 + pE[N(H)] + qE[N'(T)],$$

i.e.,

$$\beta(T) = 1 + p\beta(H) + q\beta(T).$$

## First Passage Time - Example 3

You roll a balanced six-sided die until the sum of the last two rolls is 8.  
How many times do you have to roll the die, on average?



The arrows out of  $3, \dots, 6$  (not shown) are similar to those out of 2.

$$\beta(S) = 1 + \frac{1}{6} \sum_{j=1}^6 \beta(j); \beta(1) = 1 + \frac{1}{6} \sum_{j=1}^6 \beta(j); \beta(i) = 1 + \frac{1}{6} \sum_{j=1, \dots, 6; j \neq 8-i} \beta(j), i = 2, \dots, 6.$$


Symmetry:  $\beta(2) = \dots = \beta(6) =: \gamma$ . Also,  $\beta(1) = \beta(S)$ . Thus,

$$\beta(S) = 1 + (5/6)\gamma + \beta(S)/6; \quad \gamma = 1 + (4/6)\gamma + (1/6)\beta(S).$$

$$\Rightarrow \dots \beta(S) = 8.4.$$

## First Passage Time - Example 4

You try to go up a ladder that has 20 rungs. At each time step, you succeed in going up by one rung with probability  $p = 0.9$ . Otherwise, you fall back to the ground. How many time steps does it take you to reach the top of the ladder, on average?



$$\beta(n) = 1 + p\beta(n+1) + q\beta(0), 0 \leq n < 19$$


$$\beta(19) = 1 + p\beta(20) + q\beta(0)$$

$$\Rightarrow \beta(0) = \frac{p^{-20} - 1}{1 - p} \approx 72.$$

See Lecture Note 24 for algebra.

## First Passage Time - Example 5

You play a game of “heads or tails” using a biased coin that yields ‘heads’ with probability  $p < 0.5$ . You start with \$10. At each step, if the flip yields ‘heads’, you earn \$1. Otherwise, you lose \$1. What is the probability that you reach \$100 before \$0?



Let  $\alpha(n)$  be the probability of reaching 100 before 0, starting from  $n$ , for  $n = 0, 1, \dots, 100$ .


$$\alpha(0) = 0; \alpha(100) = 1.$$

$$\alpha(n) = p\alpha(n+1) + q\alpha(n-1), 0 < n < 100.$$

$$\Rightarrow \alpha(n) = \frac{1 - \rho^n}{1 - \rho^{100}} \text{ with } \rho = qp^{-1}. \text{ (See LN 24)}$$

## First Passage Time - Example 5

You play a game of “heads or tails” using a biased coin that yields ‘heads’ with probability 0.48. You start with \$10. At each step, if the flip yields ‘heads’, you earn \$1. Otherwise, you lose \$1. What is the probability that you reach \$100 before \$0?



Morale of example: Be careful!


## Summary: First Step Equations

Let  $X_n$  be a MC on  $\mathcal{X}$  and  $A, B \subset \mathcal{X}$  with  $A \cap B = \emptyset$ . Define

$$T_A = \min\{n \geq 0 \mid X_n \in A\} \text{ and } T_B = \min\{n \geq 0 \mid X_n \in B\}.$$

Let

$$\beta(i) = E[T_A \mid X_0 = i] \text{ and } \alpha(i) = \Pr[T_A < T_B \mid X_0 = i], i \in \mathcal{X}.$$



The FSE are

$$\beta(i) = 0, i \in A$$

$$\beta(i) = 1 + \sum_j P(i, j)\beta(j), i \notin A$$

$$\alpha(i) = 1, i \in A$$

$$\alpha(i) = 0, i \in B$$

$$\alpha(i) = \sum_j P(i, j)\alpha(j), i \notin A \cup B.$$