CS-184: Computer Graphics

Lecture #16: Global Illumination

Prof. James O'Brien
University of California, Berkeley

V2008-S-16-1.0

Today

- The Rendering Equation
- Radiosity Method
- Photon Mapping
- Ambient Occlusion

2


local only
HENRIK WANN JENSEN 1999


soft shadows
HENRIK WANN JENSEN 1999


caustics
HENRIK WANN JENSEN 1999


indirect illumination
HENRIK WANN JENSEN 2000

The Rendering Equation



The light shining on x from x' is equal to:
- the emitted light from x' toward x, plus
- for each bit of surface in the scene, how much light shines from that bit onto x' and is reflected toward x, scaled appropriately

The Rendering Equation



The light shining on x from x' is equal to:
- the emitted light from x' toward x, plus
- for each bit of surface in the scene, how much light shines from that bit onto x' and is reflected toward x, scaled appropriately

$$L_s(\mathbf{x},\mathbf{x}') = \delta(\mathbf{x},\mathbf{x}')\left[E(\mathbf{x},\mathbf{x}') + \int_S \rho_{x'}(\mathbf{x},\mathbf{x}'')L_s(\mathbf{x}',\mathbf{x}'')\frac{\cos(\theta')\cos(\theta'')}{||\mathbf{x}'-\mathbf{x}''||^2}d\mathbf{x}''\right]$$

The Rendering Equation

$$L_s(\mathbf{x},\mathbf{x}') = \delta(\mathbf{x},\mathbf{x}')\left[E(\mathbf{x},\mathbf{x}') + \int_S \rho_{x'}(\mathbf{x},\mathbf{x}'')L_s(\mathbf{x}',\mathbf{x}'')\frac{\cos(\theta')\cos(\theta'')}{||\mathbf{x}'-\mathbf{x}''||^2}d\mathbf{x}''\right]$$



The Rendering Equation

$$L_s(\mathbf{x},\mathbf{x}') = \delta(\mathbf{x},\mathbf{x}')\left[E(\mathbf{x},\mathbf{x}') + \int_S \rho_{x'}(\mathbf{x},\mathbf{x}'')L_s(\mathbf{x}',\mathbf{x}'')\frac{\cos(\theta')\cos(\theta'')}{||\mathbf{x}'-\mathbf{x}''||^2}d\mathbf{x}''\right]$$

Light energy hitting x from x’



The Rendering Equation

$$L_s(\mathbf{x},\mathbf{x}') = \delta(\mathbf{x},\mathbf{x}')\left[E(\mathbf{x},\mathbf{x}') + \int_S \rho_{x'}(\mathbf{x},\mathbf{x}'')L_s(\mathbf{x}',\mathbf{x}'')\frac{\cos(\theta')\cos(\theta'')}{||\mathbf{x}'-\mathbf{x}''||^2}d\mathbf{x}''\right]$$



The Rendering Equation

$$L_s(\mathbf{x},\mathbf{x}') = \delta(\mathbf{x},\mathbf{x}')\left[E(\mathbf{x},\mathbf{x}') + \int_S \rho_{x'}(\mathbf{x},\mathbf{x}'')L_s(\mathbf{x}',\mathbf{x}'')\frac{\cos(\theta')\cos(\theta'')}{||\mathbf{x}'-\mathbf{x}''||^2}d\mathbf{x}''\right]$$

Can x see x' ?



The Rendering Equation

$$L_s(\mathbf{x},\mathbf{x}') = \delta(\mathbf{x},\mathbf{x}')\left[E(\mathbf{x},\mathbf{x}') + \int_S \rho_{x'}(\mathbf{x},\mathbf{x}'')L_s(\mathbf{x}',\mathbf{x}'')\frac{\cos(\theta')\cos(\theta'')}{||\mathbf{x}'-\mathbf{x}''||^2}d\mathbf{x}''\right]$$



The Rendering Equation

$$L_s(\mathbf{x},\mathbf{x}') = \delta(\mathbf{x},\mathbf{x}')\left[E(\mathbf{x},\mathbf{x}') + \int_S \rho_{x'}(\mathbf{x},\mathbf{x}'')L_s(\mathbf{x}',\mathbf{x}'')\frac{\cos(\theta')\cos(\theta'')}{||\mathbf{x}'-\mathbf{x}''||^2}d\mathbf{x}''\right]$$

Light emitted from x' toward x



The Rendering Equation

$$L_s(\mathbf{x},\mathbf{x}') = \delta(\mathbf{x},\mathbf{x}')\left[E(\mathbf{x},\mathbf{x}') + \int_S \rho_{x'}(\mathbf{x},\mathbf{x}'')L_s(\mathbf{x}',\mathbf{x}'')\frac{\cos(\theta')\cos(\theta'')}{||\mathbf{x}'-\mathbf{x}''||^2}d\mathbf{x}''\right]$$



The Rendering Equation

$$L_s(\mathbf{x},\mathbf{x}') = \delta(\mathbf{x},\mathbf{x}')\left[E(\mathbf{x},\mathbf{x}') + \int_S \rho_{x'}(\mathbf{x},\mathbf{x}'')L_s(\mathbf{x}',\mathbf{x}'')\frac{\cos(\theta')\cos(\theta'')}{||\mathbf{x}'-\mathbf{x}''||^2}d\mathbf{x}''\right]$$

sum over every bit of surface in the scene



The Rendering Equation

$$L_s(\mathbf{x},\mathbf{x}') = \delta(\mathbf{x},\mathbf{x}')\left[E(\mathbf{x},\mathbf{x}') + \int_S \rho_{x'}(\mathbf{x},\mathbf{x}'')L_s(\mathbf{x}',\mathbf{x}'')\frac{\cos(\theta')\cos(\theta'')}{||\mathbf{x}'-\mathbf{x}''||^2}d\mathbf{x}''\right]$$



The Rendering Equation

$$L_s(\mathbf{x},\mathbf{x}') = \delta(\mathbf{x},\mathbf{x}')\left[E(\mathbf{x},\mathbf{x}') + \int_S \rho_{x'}(\mathbf{x},\mathbf{x}'')L_s(\mathbf{x}',\mathbf{x}'')\frac{\cos(\theta')\cos(\theta'')}{||\mathbf{x}'-\mathbf{x}''||^2}d\mathbf{x}''\right]$$

Light emitted from x” toward x’



The Rendering Equation

$$L_s(\mathbf{x},\mathbf{x}') = \delta(\mathbf{x},\mathbf{x}')\left[E(\mathbf{x},\mathbf{x}') + \int_S \rho_{x'}(\mathbf{x},\mathbf{x}'')L_s(\mathbf{x}',\mathbf{x}'')\frac{\cos(\theta')\cos(\theta'')}{||\mathbf{x}'-\mathbf{x}''||^2}d\mathbf{x}''\right]$$



The Rendering Equation

$$L_s(\mathbf{x},\mathbf{x}') = \delta(\mathbf{x},\mathbf{x}')\left[E(\mathbf{x},\mathbf{x}') + \int_S \rho_{x'}(\mathbf{x},\mathbf{x}'')L_s(\mathbf{x}',\mathbf{x}'')\frac{\cos(\theta')\cos(\theta'')}{||\mathbf{x}'-\mathbf{x}''||^2}d\mathbf{x}''\right]$$

scaled down by the BRDF of x'



The Rendering Equation

$$L_s(\mathbf{x},\mathbf{x}') = \delta(\mathbf{x},\mathbf{x}')\left[E(\mathbf{x},\mathbf{x}') + \int_S \rho_{x'}(\mathbf{x},\mathbf{x}'')L_s(\mathbf{x}',\mathbf{x}'')\frac{\cos(\theta')\cos(\theta'')}{||\mathbf{x}'-\mathbf{x}''||^2}d\mathbf{x}''\right]$$



The Rendering Equation

$$L_s(\mathbf{x},\mathbf{x}') = \delta(\mathbf{x},\mathbf{x}')\left[E(\mathbf{x},\mathbf{x}') + \int_S \rho_{x'}(\mathbf{x},\mathbf{x}'')L_s(\mathbf{x}',\mathbf{x}'')\frac{\cos(\theta')\cos(\theta'')}{||\mathbf{x}'-\mathbf{x}''||^2}d\mathbf{x}''\right]$$

scaled down by distance and relative orientation ("form factor")



The Rendering Equation

$$L_s(\mathbf{x},\mathbf{x}') = \delta(\mathbf{x},\mathbf{x}')\left[E(\mathbf{x},\mathbf{x}') + \int_S \rho_{x'}(\mathbf{x},\mathbf{x}'')L_s(\mathbf{x}',\mathbf{x}'')\frac{\cos(\theta')\cos(\theta'')}{||\mathbf{x}'-\mathbf{x}''||^2}d\mathbf{x}''\right]$$



Radiosity

- Assume all materials are perfectly Lambertian (diffuse only, no specularities)
  - Removes all dependance on directions
  - Reduces dimensionality of lightfield
  - Allows a FEM solution (break up into chunks)
- Can also relax assumption slightly...

6

Early radiosity

from Hanrahan 2000

Assume Lambertian

$$L_s(\mathbf{x},\mathbf{x}') = \delta(\mathbf{x},\mathbf{x}')\left[E(\mathbf{x},\mathbf{x}') + \int_S \rho_{x'}(\mathbf{x},\mathbf{x}'')L_s(\mathbf{x}',\mathbf{x}'')\frac{\cos(\theta')\cos(\theta'')}{||\mathbf{x}'-\mathbf{x}''||^2}\mathrm{d}\mathbf{x}''\right]$$

$$L_s(\mathbf{x},\mathbf{x}') = \delta(\mathbf{x},\mathbf{x}')\left[E_{x'} + \int_S \rho_{x'}L_s(\mathbf{x}',\mathbf{x}'')\frac{\cos(\theta')\cos(\theta'')}{||\mathbf{x}'-\mathbf{x}''||^2}\mathrm{d}\mathbf{x}''\right]$$



Assume Lambertian

$$L_s(\mathbf{x},\mathbf{x}') = \delta(\mathbf{x},\mathbf{x}')\left[E(\mathbf{x},\mathbf{x}') + \int_S \rho_{x'}(\mathbf{x},\mathbf{x}'')L_s(\mathbf{x}',\mathbf{x}'')\frac{\cos(\theta')\cos(\theta'')}{||\mathbf{x}'-\mathbf{x}''||^2}\mathrm{d}\mathbf{x}''\right]$$

$$L_s(\mathbf{x},\mathbf{x}') = \underline{\delta(\mathbf{x},\mathbf{x}')}\left[E_{x'} + \int_S \rho_{x'}L_s(\mathbf{x}',\mathbf{x}'')\frac{\cos(\theta')\cos(\theta'')}{||\mathbf{x}'-\mathbf{x}''||^2}\mathrm{d}\mathbf{x}''\right]$$

Only term dependent on $\mathbf{x}$



Rewrite in Terms of Radiosity

$$L_s(\mathbf{x},\mathbf{x}') = \delta(\mathbf{x},\mathbf{x}')\left[E_{x'} + \int_S \rho_{x'} L_s(\mathbf{x}',\mathbf{x}'')\frac{\cos(\theta')\cos(\theta'')}{||\mathbf{x}'-\mathbf{x}''||^2}\mathrm{d}\mathbf{x}''\right]$$

$$H_{x'} = E_{x'} + \rho_{x'}\int_S \delta(\mathbf{x}',\mathbf{x}'')\frac{H_{x''}}{2\pi}\frac{\cos(\theta')\cos(\theta'')}{||\mathbf{x}'-\mathbf{x}''||^2}\mathrm{d}\mathbf{x}''$$



Note: we changed defn of E here.

Discretize into Patches





Piece-wise constant patches



Example mesh for Cornell Box
by Mark Schmelzenbach

10

Discretize into Patches



The Candlestick Theater,
Mark Mack Architects.

Discretize into Patches

The Candlestick Theater,
Mark Mack Architects.

12

Rewrite in Terms of Patches

$$H_{x'} = E_{x'} + \rho_{x'} \int_S \delta(\mathbf{x}', \mathbf{x}'') \frac{H_{x''} \cos(\theta') \cos(\theta'')}{||\mathbf{x}' - \mathbf{x}''||^2} d\mathbf{x}''$$

$$H_i = E_i + \rho_i \sum_j H_j \int_{S_j} \delta_{ij} \frac{\cos(\theta_i) \cos(\theta_j)}{2\pi ||\mathbf{c}_i - \mathbf{x}||^2} d\mathbf{x}$$



13

Rewrite in Terms of Patches

$$H_{x'} = E_{x'} + \rho_{x'} \int_S \delta(\mathbf{x}', \mathbf{x}'') \frac{H_{x''} \cos(\theta') \cos(\theta'')}{2\pi \quad ||\mathbf{x}' - \mathbf{x}''||^2} d\mathbf{x}''$$

$$H_i = E_i + \rho_i \sum_j H_j \int_{S_j} \delta_{ij} \frac{\cos(\theta_i) \cos(\theta_j)}{2\pi ||\mathbf{c}_i - \mathbf{x}||^2} d\mathbf{x}$$

Form factor from j to i, F_{ij}



13

Rewrite in Terms of Patches

$$H_{x'} = E_{x'} + \rho_{x'} \int_S \delta(\mathbf{x}', \mathbf{x}'') \frac{H_{x''} \cos(\theta') \cos(\theta'')}{||\mathbf{x}' - \mathbf{x}''||^2} d\mathbf{x}''$$

$$H_i = E_i + \rho_i \sum_j H_j \int_{S_j} \delta_{ij} \frac{\cos(\theta_i) \cos(\theta_j)}{2\pi ||\mathbf{c}_i - \mathbf{x}||^2} d\mathbf{x}$$

Form factor from j to i, F_{ij}



Example of a rough approximation:

$$F_{ij} \approx \delta_{ij} \frac{\cos(\theta_i) \cos(\theta_j)}{2\pi ||\mathbf{c}_i - \mathbf{c}_j||^2} A_j$$

13

Radiosity Method

- Given the E_i and ρ_i
- First compute F_{ij}
- Then solve $H_i = E_i + \rho_i \sum_j H_j F_{ij}$

$$\mathbf{h} = \mathbf{e} + \mathbf{A}\mathbf{h}$$
$$(\mathbf{I} - \mathbf{A})\mathbf{h} = \mathbf{e}$$

- Comments:
  - The matrix $\mathbf{A}$ is typically very large
  - It is also sparse (why?)
  - Should be solved with an iterative method
    - e.g.: Jacobi or Gauss-Seidel
  - **Solution is view independent**

14

Radiosity Method

- Given the light emitted and surface properties
- First compute F_{ij}, form factors between patches
- Then **solve a linear system to balance energy between all patches**
- Comments:
  - The system is very large
  - It is also sparse (why?)
  - Should be solved with an iterative method
    - e.g.: Jacobi or Gauss-Seidel
  - **Solution is view independent**

15

Progressive Radiosity

- If magnitude of eigenvalues of $\mathbf{A}$<1

$$(\mathbf{I}-\mathbf{A})^{-1}=\mathbf{I}+\mathbf{A}+\mathbf{A}^2+\mathbf{A}^3+\cdots$$

  - True for form-factor matrices
  - $\mathbf{h}^{k+1}=\mathbf{h}^k+\mathbf{u}^{k+1}$
    $\mathbf{u}^{k+1}=\mathbf{A}\,\mathbf{u}^k$
    $\mathbf{h}^0=0 \quad \mathbf{u}^0=\mathbf{e}$

Idea: let important sources of light energy emit first, maybe don't even bother with dark things

  - Use Gauss-Seidel-like iteration but reorder by priority

Southwell Relaxation

16

Progressive Radiosity



From dissertation "Efficient and predictive realistic image synthesis"
by Karol Myszkowski

17

Touchup

- Each patch will have a constant color
  - Smooth solution (e.g. average to vertices)

Example mesh for Cornell Box
by Mark Schmelzenbach

Does not match but you get the idea...

18

Other Things

- Each patch will have a constant color
  - Smooth solution (e.g. average to vertices)
- No specular reflection
  - Add Phong specular term or raytraced specular reflection
- Grid artifacts
  - Be clever with grid...

19

Hierarchical Radiosity

- Light smoothes with distance
  - Compare $1/h^2$ with $1/(h^2+d^2)$ as h gets large



Hierarchical Radiosity

- Light smoothes with distance
  - Compare $1/h^2$ with $1/(h^2+d^2)$ as h gets large
- Group patches into hierarchy
  - Far interactions use lower-res form factors



Computing Form Factors

- Form factors have a geometric meaning





Images from
SIGGRAPH 93 Education Slide Set
by Stephen Spencer

22

Computing Form Factors

- Form factors have a geometric meaning
- "Hemicube" algorithm uses regular scan conversion



Images from
SIGGRAPH 93 Education Slide Set
by Stephen Spencer

23

Computing Form Factors

- Form factors have a geometric meaning
- "Hemicube" algorithm uses regular scan conversion
- Also computed by ray-based sampling
- **In practice, computing form factors is the bottleneck**

24

Photon Mapping

- Lights cast "photons" into environment
  - Cast in random directions
  - Trace into environment
  - Store records at intersections



25

Photon Mapping

- Lights cast "photons" into environment
  - Cast in random directions
  - Trace into environment
  - Store records at intersections
    - With KD-Trees...



26

Comparison



Ray Tracing



Ray Tracing w/ Photon Map

Catherine Bendebury and Jonathan Michaels
CS 184 Spring 2005

27

Photon Mapping

Image by Per Christensen

A ray traced image

Note:
- Dark shadows
- Unlit corners
- Nice reflections

28

Photon Mapping



Image by Per Christensen

Raw photons

Note:
- Noisy
- Sparse

29

Photon Mapping

Image by Per Christensen

Interpolated Photons

Note:
- Still noisy
- Biased

30

Photon Mapping

Image by Per Christensen

Interpolated Photons
(multiplied by diffuse)

Note:
- Still noisy
- Biased

31

Photon Mapping

- Final Gather
  - Ray trace scene
  - Direct and specular rays as normal
  - Diffuse rays traced into photon map
  - *Diffuse reflection smoothes noise*

32

Photon Mapping

Image by Per Christensen

Final Image

Note:
- Not noisy
- Nice lighting
- Reflections
- May still be biased

Final gather often bottleneck...

33

Ambient Occlusion

- A "hack" to create more realistic ambient illumination cheaply
- Assume light from everywhere is partially blocked by local objects
  - At a point on the surface cast rays at random
  - Ambient term is proportional to percent of rays that hit nothing
  - Weight average by cosine of angle with normal
  - Take into account how far before occluded

34

Ambient Occlusion



35

Ambient Occlusion

Diffuse Only

Ambient Occlusion

Combined

36

Ambient Occlusion



nVidia Gelato Demo Image

37