High Dynamic Range Images

© Alyosha Efros

CS194: Image Manipulation & Computational Photography

…with a lot of slides stolen from Paul Debevec

Alexei Efros, UC Berkeley, Fall 2017

Why HDR?



Problem: Dynamic Range



1

1500

25,000

400,000

2,000,000,000

The real world is high dynamic range.

Image



pixel (312, 284) = **42**

42 photos?

Long Exposure



Short Exposure



Camera Calibration

- Geometric
  - How pixel **coordinates** relate to **directions** in the world
- Photometric
  - How pixel **values** relate to **radiance** amounts in the world



The Image Acquisition Pipeline


Development
CCD
ADC
Remapping
film density
analog voltages
digital values
pixel values

Imaging system response function



log Exposure = log (Radiance * Δt)

(CCD photon count)

Varying Exposure

Camera is not a photometer!

- Limited dynamic range
  - $\Rightarrow$ Perhaps use multiple exposures?
- Unknown, nonlinear response
  - $\Rightarrow$ Not possible to convert pixel values to radiance
- Solution:
  - Recover response curve from multiple exposures, then reconstruct the ***radiance map***

Recovering High Dynamic Range Radiance Maps from Photographs

Paul Debevec
Jitendra Malik

Computer Science Division
University of California at Berkeley

August 1997

Ways to vary exposure

Ways to vary exposure

- Shutter Speed (*)
- F/stop (aperture, iris)

- Neutral Density (ND) Filters

Shutter Speed

- **Ranges: Canon D30: 30 to 1/4,000 sec.**
- **Sony VX2000: ¼ to 1/10,000 sec.**
- **Pros:**
- **Directly varies the exposure**
- **Usually accurate and repeatable**
- **Issues:**
- **Noise in long exposures**

Shutter Speed

- **Note: shutter times usually obey a power series – each “stop” is a factor of 2**
- **¼, 1/8, 1/15, 1/30, 1/60, 1/125, 1/250, 1/500, 1/1000 sec**
- **Usually really is:**
- **¼, 1/8, 1/16, 1/32, 1/64, 1/128, 1/256, 1/512, 1/1024 sec**

The Algorithm

Image series



$\Delta t =$	$\Delta t =$	$\Delta t =$	$\Delta t =$	$\Delta t =$
1/64 sec	**1/16 sec**	**1/4 sec**	**1 sec**	**4 sec**

Pixel Value Z = f(Exposure)

Exposure = Radiance ´ Δt

log Exposure = log Radiance + log Δt

Response Curve

Assuming unit radiance for each pixel



After adjusting radiances to obtain a smooth response



The Math

- Let *g(z)* be the *discrete* inverse response function
- For each pixel site *i* in each image *j*, want:

$$\ln Radiance_i + \ln \Delta t_j = g(Z_{ij})$$

- Solve the overdetermined linear system:

$$\underbrace{\sum_{i=1}^{N}\sum_{j=1}^{P}\left[\ln Radiance_i + \ln \Delta t_j - g(Z_{ij})\right]^2}_{\text{fitting term}} + \underbrace{\lambda \sum_{z=Z_{min}}^{Z_{max}} g''(z)^2}_{\text{smoothness term}}$$

Matlab Code

```
function [g,lE]=gsolve(Z,B,l,w)

n = 256;
A = zeros(size(Z,1)*size(Z,2)+n+1,n+size(Z,1));
b = zeros(size(A,1),1);

k = 1;                  %% Include the data-fitting equations
for i=1:size(Z,1)
  for j=1:size(Z,2)
    wij = w(Z(i,j)+1);
    A(k,Z(i,j)+1) = wij; A(k,n+i) = -wij; b(k,1) = wij * B(i,j);
    k=k+1;
  end
end

A(k,129) = 1;           %% Fix the curve by setting its middle value to
k=k+1;

for i=1:n-2             %% Include the smoothness equations
  A(k,i)=l*w(i+1); A(k,i+1)=-2*l*w(i+1); A(k,i+2)=l*w(i+1);
  k=k+1;
end

x = A\b;                %% Solve the system using SVD

g = x(1:n);
lE = x(n+1:size(x,1));
```

Results: Digital Camera

Kodak DCS460
1/30 to 30 sec



Recovered response curve



Reconstructed radiance map

Results: Color Film

- Kodak Gold ASA 100, PhotoCD



Recovered Response Curves



The Radiance Map



The Radiance Map



Linearly scaled to display device

Now What?



Tone Mapping

- How can we do this?

  Linear scaling?, thresholding? Suggestions?



Simple Global Operator

- Compression curve needs to
  - Bring everything within range
  - Leave dark areas alone
- In other words
  - Asymptote at 255
  - Derivative of 1 at 0

Global Operator (Reinhart et al)

$$L_{display} = \frac{L_{world}}{1 + L_{world}}$$



Global Operator Results



Reinhart Operator



Darkest 0.1% scaled to display device

What do *we* see?



Vs.



What does the eye sees?



Figure 1: The range of luminances in the natural environment and associated visual parameters. After Hood (1986).

The eye has a huge dynamic range
Do we see a true radiance map?

Metamores





Can we use this for range compression?

Compressing Dynamic Range