

61A Extra Lecture 1


Announcements

- If you want 1 unit (pass/no pass) of credit for this CS 98, stay tuned for a Piazza post
 - Only for people who really want extra work that's beyond the scope of normal CS 61A
- Anyone is welcome to attend the extra lectures, whether or not they enroll
- Permanent lecture times are TBD, but probably Wednesday evening or Monday evening

Newton's Method

Newton's Method Background

Quickly finds accurate approximations to zeroes of differentiable functions!



Application: a method for computing square roots, cube roots, etc.

The positive zero of $f(x) = x^2 - a$ is \sqrt{a} . (We're solving the equation $x^2 = a$.)

Newton's Method

Given a function f and initial guess x ,

Repeatedly improve x :


Compute the value of f
at the guess: $f(x)$

Compute the derivative
of f at the guess: $f'(x)$

Update guess x to be:


$$x - \frac{f(x)}{f'(x)}$$

Finish when $f(x) = 0$ (or close enough)



Using Newton's Method

How to find the square root of 2?




```
>>> f = lambda x: x*x - 2
>>> df = lambda x: 2*x
>>> find_zero(f, df)
1.4142135623730951
```

$$f(x) = x^2 - 2$$
$$f'(x) = 2x$$

Applies Newton's method

How to find the cube root of 729?



```
>>> g = lambda x: x*x*x - 729
>>> dg = lambda x: 3*x*x
>>> find_zero(g, dg)
9.0
```

$$g(x) = x^3 - 729$$
$$g'(x) = 3x^2$$

Iterative Improvement

Special Case: Square Roots

How to compute `square_root(a)`

Idea: Iteratively refine a guess x about the square root of a

Update:
$$x = \frac{x + \frac{a}{x}}{2}$$

(Demo)

Babylonian Method

Implementation questions:

What guess should start the computation?

How do we know when we are finished?

Special Case: Cube Roots

How to compute `cube_root(a)`

Idea: Iteratively refine a guess `x` about the cube root of `a`

Update:
$$x = \frac{2 \cdot x + \frac{a}{x^2}}{3}$$

(Demo)

Implementation questions:

What guess should start the computation?

How do we know when we are finished?

Implementing Newton's Method

(Demo)

Extensions

Approximate Differentiation

Differentiation can be performed symbolically or numerically

$$f(x) = x^2 - 16$$


$$f'(x) = 2x$$

$$f'(2) = 4$$

$$f'(x) = \lim_{a \rightarrow 0} \frac{f(x+a) - f(x)}{a}$$

$$f'(x) \approx \frac{f(x+a) - f(x)}{a} \quad (\text{if } a \text{ is small})$$

(Demo)



Critical Points and Inverses

Maxima, minima, and inflection points of a differentiable function occur when the derivative is 0

(Demo)

The inverse $f^{-1}(y)$ of a differentiable, one-to-one function computes the value x such that $f(x) = y$

(Demo)

