

CS-184: Computer Graphics

Lecture #6: Raytracing

Prof. James O'Brien
University of California, Berkeley

V2009-F-06-1.0

Today

- Raytracing
 - Shadows and direct lighting
 - Reflection and refraction
 - Antialiasing, motion blur, soft shadows, and depth of field
- Intersection Tests
 - Ray-primitive

Raytracing Assignment



3

Light in an Environment



Lady writing a Letter with her Maid
National Gallery of Ireland, Dublin
Johannes Vermeer, 1670

4

Global Illumination Effects



PCKTWCH
Kevin Odhner
POV-Ray

5


Global Illumination Effects



A Philco 6Z4 Vacuum Tube
Steve Anger
POV-Ray

6


Global Illumination Effects



Caustic Sphere
Henrik Jensen
(refraction caustic)

7


Global Illumination Effects



Caustic Ring
Henrik Jensen
(reflection caustic)

8

Global Illumination Effects



Sphere Flake
Henrik Jensen

9

Early Raytracing



Turner Whitted


10

Raytracing

- Scan conversion
 - $3D \rightarrow 2D \rightarrow \text{Image}$
 - Based on transforming geometry
- Raytracing
 - $3D \rightarrow \text{Image}$
 - Geometric reasoning about light rays

11


Raytracing



Eye, view plane section, and scene

12


Raytracing



Launch ray from eye through pixel, see what it hits

13

Raytracing



Compute color and fill-in the pixel

14


Raytracing

- Basic tasks
 - Build a ray
 - Figure out what a ray hits
 - Compute shading

15

Building Eye Rays

- Rectilinear image plane build from four points




$$\mathbf{P} = u (v\mathbf{LL} + (1 - v)\mathbf{UL}) + (1 - u)(v\mathbf{LR} + (1 - v)\mathbf{UR})$$

16


Building Eye Rays

- Nonlinear projections
 - Non-planar projection surface
 - Variable eye location



17


Examples



Multiple-Center-of-Projection Images
P. Rademacher and G. Bishop
SIGGRAPH 1998

18

Examples



Spherical and Cylindrical Projections
Ben Kreunen
From *Big Ben's Panorama Tutorials*

19


Building Eye Rays

- Ray equation

$$\mathbf{R}(t) = \mathbf{E} + t(\mathbf{P} - \mathbf{E})$$

$$t \in [1 \dots + \infty]$$

- Through eye at $t = 0$
- At pixel center at $t = 1$




20

Shadow Rays

- Detect shadow by rays to light source

$$\mathbf{R}(t) = \mathbf{S} + t(\mathbf{L} - \mathbf{S})$$

$$t \in [\epsilon \dots 1)$$




21

Shadow Rays

- Test for occluder
 - No occluder, shade normally (e.g. Phong model)
 - Yes occluder, skip light (don't skip ambient)
- Self shadowing
 - Add shadow bias
 - Test object ID



Self-shadowing



Correct

22


Reflection Rays

- Recursive shading

- Ray bounces off object
- Treat bounce rays (mostly) like eye rays
- Shade bounce ray and return color
 - Shadow rays
 - Recursive reflections
- Add color to shading at original point
 - Specular or separate reflection coefficient

$$\mathbf{R}(t) = \mathbf{S} + t\mathbf{B}$$

$$t \in [\epsilon \dots +\infty)$$





23

Reflection Rays

- Recursion Depth

- Truncate at fixed number of bounces
- Multiplier less than J.N.D.




24

Refracted Rays

- Transparent materials bend light

- Snell's Law $\frac{n_i}{n_t} = \frac{\sin \theta_t}{\sin \theta_i}$ (see clever formula in text...)

$\sin \theta_t > 1$ Total (internal) reflection



Refracted Rays

- Coefficient on transmitted ray depends on θ


- Schlick approximation to Fresnel Equations

$$k_t(\theta_i) = k_0 + (1 - k_0)(1 - \cos \theta_i)^5$$

$$k_0 = \left(\frac{n_t - 1}{n_t + 1} \right)^2$$


- Attenuation
 - Wavelength (color) dependant
 - Exponential with distance

Refracted Rays



Anti-Aliasing



- Boolean on/off for pixels causes problems
 - Consider scan conversion algorithm:



- Compute z through each pixel center
- Recall Nyquist Theorem
 - *Sampling rate \geq twice highest frequency*

Anti-Aliasing




- Desired solution of an integral over pixel



29

“Distributed” Raytracing

- Send multiple rays through each pixel




One Sample 5x5 Grid 5x5 Jittered Grid


- Average results together
- Jittering trades aliasing for noise

30

“Distributed” Raytracing



31




Beverly Chiu and Max Delgado
CS 184 2007


“Distributed” Raytracing

- Use multiple rays for reflection and refraction
 - At each bounce send out many extra rays
 - Quasi-random directions
 - Use BRDF (or Phong approximation) for weights
- How many rays?


33



|



16



256

Soft Shadows





Figure from S. Cheney

Soft Shadows






Figure from S. Cheney

Soft Shadows

- Distribute shadow rays over light surface




All shadow rays
go through

No shadow rays
go through

Some shadow
rays go through

Figure from S. Cheney ³⁸



Motion Blur


- Distribute rays over *time*
 - More when we talk about animation...



Pool Balls
Tom Porter
RenderMan

40


Depth of Field




Kolb, Mitchell, and Hanrahan
SIGGRAPH 1995

41


Depth of Field



No DoF




Jittered rays for DoF



More rays





Multiple images for DoF



Even more rays


42

Other Lens Effects




Ray -vs- Sphere Test

- Ray equation: $\mathbf{R}(t) = \mathbf{A} + t\mathbf{D}$
- Implicit equation for sphere: $|\mathbf{X} - \mathbf{C}|^2 - r^2 = 0$
- Combine:
$$|\mathbf{R}(t) - \mathbf{C}|^2 - r^2 = 0$$
$$|\mathbf{A} + t\mathbf{D} - \mathbf{C}|^2 - r^2 = 0$$
- Quadratic equation in t



44

Ray -vs- Sphere Test



45

Ray -vs- Triangle

• Ray equation: $\mathbf{R}(t) = \mathbf{A} + t\mathbf{D}$

• Triangle in barycentric coordinates:

$$\mathbf{X}(\beta, \gamma) = \mathbf{V}_1 + \beta(\mathbf{V}_2 - \mathbf{V}_1) + \gamma(\mathbf{V}_3 - \mathbf{V}_1)$$

• Combine:

$$\mathbf{V}_1 + \beta(\mathbf{V}_2 - \mathbf{V}_1) + \gamma(\mathbf{V}_3 - \mathbf{V}_1) = \mathbf{A} + t\mathbf{D}$$

• Solve for β , γ , and t

- 3 equations 3 unknowns
- Beware divide by near-zero
- Check ranges

