


CS 188: Artificial Intelligence

Bayes' Nets




Instructors: Dan Klein and Pieter Abbeel --- University of California, Berkeley

Probabilistic Models

- Models describe how (a portion of) the world works
- **Models are always simplifications**
 - May not account for every variable
 - May not account for all interactions between variables
 - “All models are wrong; but some are useful.”
– George E. P. Box
- What do we do with probabilistic models?
 - We (or our agents) need to reason about unknown variables, given evidence
 - Example: explanation (diagnostic reasoning)
 - Example: prediction (causal reasoning)
 - Example: value of information



Independence



Independence


- Two variables are *independent* if:

$$\forall x, y : P(x, y) = P(x)P(y)$$

- This says that their joint distribution *factors* into a product two simpler distributions
- Another form:

$$\forall x, y : P(x|y) = P(x)$$

- We write: $X \perp\!\!\!\perp Y$
- Independence is a simplifying *modeling assumption*
 - Empirical* joint distributions: at best “close” to independent
 - What could we assume for {Weather, Traffic, Cavity, Toothache}?



Example: Independence?

$P_1(T, W)$

| T | W | P |
|------|------|-----|
| hot | sun | 0.4 |
| hot | rain | 0.1 |
| cold | sun | 0.2 |
| cold | rain | 0.3 |

$P(T)$

| T | P |
|------|-----|
| hot | 0.5 |
| cold | 0.5 |

$P_2(T, W)$

| T | W | P |
|------|------|-----|
| hot | sun | 0.3 |
| hot | rain | 0.2 |
| cold | sun | 0.3 |
| cold | rain | 0.2 |

$P(W)$

| W | P |
|------|-----|
| sun | 0.6 |
| rain | 0.4 |

Example: Independence

- N fair, independent coin flips:

$P(X_1)$

| | |
|---|-----|
| H | 0.5 |
| T | 0.5 |


$P(X_2)$

| | |
|---|-----|
| H | 0.5 |
| T | 0.5 |

...


$P(X_n)$

| | |
|---|-----|
| H | 0.5 |
| T | 0.5 |




$P(X_1, X_2, \dots, X_n)$

2^n



Conditional Independence

- $P(\text{Toothache}, \text{Cavity}, \text{Catch})$
- If I have a cavity, the probability that the probe catches in it doesn't depend on whether I have a toothache:
 - $P(+\text{catch} \mid +\text{toothache}, +\text{cavity}) = P(+\text{catch} \mid +\text{cavity})$
- The same independence holds if I don't have a cavity:
 - $P(+\text{catch} \mid +\text{toothache}, -\text{cavity}) = P(+\text{catch} \mid -\text{cavity})$
- Catch is *conditionally independent* of Toothache given Cavity:
 - $P(\text{Catch} \mid \text{Toothache}, \text{Cavity}) = P(\text{Catch} \mid \text{Cavity})$
- Equivalent statements:
 - $P(\text{Toothache} \mid \text{Catch}, \text{Cavity}) = P(\text{Toothache} \mid \text{Cavity})$
 - $P(\text{Toothache}, \text{Catch} \mid \text{Cavity}) = P(\text{Toothache} \mid \text{Cavity}) P(\text{Catch} \mid \text{Cavity})$
 - One can be derived from the other easily



Conditional Independence

- Unconditional (absolute) independence very rare (why?)
- *Conditional independence* is our most basic and robust form of knowledge about uncertain environments.

- X is conditionally independent of Y given Z

$$X \perp\!\!\!\perp Y | Z$$

if and only if:

$$\forall x, y, z : P(x, y | z) = P(x | z)P(y | z)$$

or, equivalently, if and only if

$$\forall x, y, z : P(x | z, y) = P(x | z)$$



Conditional Independence

- What about this domain:
 - Traffic
 - Umbrella
 - Raining



Conditional Independence

- What about this domain:
 - Fire
 - Smoke
 - Alarm



Conditional Independence and the Chain Rule

- Chain rule: $P(X_1, X_2, \dots, X_n) = P(X_1)P(X_2|X_1)P(X_3|X_1, X_2) \dots$


- Trivial decomposition:

$$P(\text{Traffic, Rain, Umbrella}) = P(\text{Rain})P(\text{Traffic}|\text{Rain})P(\text{Umbrella}|\text{Rain, Traffic})$$

- With assumption of conditional independence:

$$P(\text{Traffic, Rain, Umbrella}) = P(\text{Rain})P(\text{Traffic}|\text{Rain})P(\text{Umbrella}|\text{Rain})$$

- Bayes' nets / graphical models help us express conditional independence assumptions



Ghostbusters Chain Rule

- Each sensor depends only on where the ghost is
- That means, the two sensors are conditionally independent, given the ghost position
- T: Top square is red
B: Bottom square is red
G: Ghost is in the top




- Givens:
 $P(+g) = 0.5$
 $P(-g) = 0.5$
 $P(+t \mid +g) = 0.8$
 $P(+t \mid -g) = 0.4$
 $P(+b \mid +g) = 0.4$
 $P(+b \mid -g) = 0.8$

$$P(T,B,G) = P(G) P(T \mid G) P(B \mid G)$$

| T | B | G | P(T,B,G) |
|----|----|----|----------|
| +t | +b | +g | 0.16 |
| +t | +b | -g | 0.16 |
| +t | -b | +g | 0.24 |
| +t | -b | -g | 0.04 |
| -t | +b | +g | 0.04 |
| -t | +b | -g | 0.24 |
| -t | -b | +g | 0.06 |
| -t | -b | -g | 0.06 |




Bayes' Nets: Big Picture




Bayes' Nets: Big Picture


- Two problems with using full joint distribution tables as our probabilistic models:
 - Unless there are only a few variables, the joint is WAY too big to represent explicitly
 - Hard to learn (estimate) anything empirically about more than a few variables at a time
- **Bayes' nets**: a technique for describing complex joint distributions (models) using simple, local distributions (conditional probabilities)
 - More properly called **graphical models**
 - We describe how variables locally interact
 - Local interactions chain together to give global, indirect interactions
 - For about 10 min, we'll be vague about how these interactions are specified



Example Bayes' Net: Insurance





Example Bayes' Net: Car





Graphical Model Notation

- **Nodes: variables (with domains)**
 - Can be assigned (observed) or unassigned (unobserved)
- **Arcs: interactions**
 - Similar to CSP constraints
 - Indicate “direct influence” between variables
 - Formally: encode conditional independence (more later)
- For now: imagine that arrows mean direct causation (in general, they don't!)



Example: Coin Flips

- N independent coin flips



- No interactions between variables: **absolute independence**



Example: Traffic

- Variables:




- R: It rains
- T: There is traffic



- Model 1: independence



- Model 2: rain causes traffic



- Why is an agent using model 2 better?

Example: Traffic II


- Let's build a causal graphical model!
- Variables
 - T: Traffic
 - R: It rains
 - L: Low pressure
 - D: Roof drips
 - B: Ballgame
 - C: Cavity



Example: Alarm Network

- Variables

- B: Burglary
- A: Alarm goes off
- M: Mary calls
- J: John calls
- E: Earthquake!



Bayes' Net Semantics




Bayes' Net Semantics



- A set of nodes, one per variable X
- A directed, acyclic graph
- A conditional distribution for each node
 - A collection of distributions over X , one for each combination of parents' values

$$P(X|a_1 \dots a_n)$$

- CPT: conditional probability table
- Description of a noisy “causal” process



A Bayes net = Topology (graph) + Local Conditional Probabilities



Probabilities in BNs



- Bayes' nets **implicitly** encode joint distributions
 - As a product of local conditional distributions
 - To see what probability a BN gives to a full assignment, multiply all the relevant conditionals together:

$$P(x_1, x_2, \dots, x_n) = \prod_{i=1}^n P(x_i | \text{parents}(X_i))$$

- Example:



$$P(+cavity, +catch, -toothache)$$

Probabilities in BNs



- Why are we guaranteed that setting

$$P(x_1, x_2, \dots, x_n) = \prod_{i=1}^n P(x_i | \text{parents}(X_i))$$

results in a proper joint distribution?

- Chain rule (valid for all distributions): $P(x_1, x_2, \dots, x_n) = \prod_{i=1}^n P(x_i | x_1 \dots x_{i-1})$



- Assume conditional independences: $P(x_i | x_1, \dots, x_{i-1}) = P(x_i | \text{parents}(X_i))$

→ Consequence: $P(x_1, x_2, \dots, x_n) = \prod_{i=1}^n P(x_i | \text{parents}(X_i))$

- Not every BN can represent every joint distribution

- The topology enforces certain conditional independencies


Example: Coin Flips



$$P(h, h, t, h) =$$

Only distributions whose variables are absolutely independent can be represented by a Bayes' net with no arcs.

Example: Traffic



$$P(R)$$

| | |
|----|-----|
| +r | 1/4 |
| -r | 3/4 |

$$P(+r, -t) =$$


$$P(T|R)$$

| | | |
|----|----|-----|
| +r | +t | 3/4 |
| +r | -t | 1/4 |
| -r | +t | 1/2 |
| -r | -t | 1/2 |




Example: Alarm Network

| B | P(B) |
|----|-------|
| +b | 0.001 |
| -b | 0.999 |



| E | P(E) |
|----|-------|
| +e | 0.002 |
| -e | 0.998 |




| A | J | P(J A) |
|----|----|--------|
| +a | +j | 0.9 |
| +a | -j | 0.1 |
| -a | +j | 0.05 |
| -a | -j | 0.95 |

| A | M | P(M A) |
|----|----|--------|
| +a | +m | 0.7 |
| +a | -m | 0.3 |
| -a | +m | 0.01 |
| -a | -m | 0.99 |

| B | E | A | P(A B,E) |
|----|----|----|----------|
| +b | +e | +a | 0.95 |
| +b | +e | -a | 0.05 |
| +b | -e | +a | 0.94 |
| +b | -e | -a | 0.06 |
| -b | +e | +a | 0.29 |
| -b | +e | -a | 0.71 |
| -b | -e | +a | 0.001 |
| -b | -e | -a | 0.999 |

Example: Traffic

- Causal direction



$P(R)$

| | |
|----|-----|
| +r | 1/4 |
| -r | 3/4 |


$P(T|R)$

| | | |
|----|----|-----|
| +r | +t | 3/4 |
| | -t | 1/4 |

| | | |
|----|----|-----|
| -r | +t | 1/2 |
| | -t | 1/2 |


$P(T, R)$

| | | |
|----|----|------|
| +r | +t | 3/16 |
| +r | -t | 1/16 |
| -r | +t | 6/16 |
| -r | -t | 6/16 |



Example: Reverse Traffic

- Reverse causality?



$P(T)$

| | |
|----|------|
| +t | 9/16 |
| -t | 7/16 |

$P(R|T)$

| | | |
|----|----|-----|
| +t | +r | 1/3 |
| | -r | 2/3 |

| | | |
|----|----|-----|
| -t | +r | 1/7 |
| | -r | 6/7 |




$P(T, R)$

| | | |
|----|----|------|
| +r | +t | 3/16 |
| +r | -t | 1/16 |
| -r | +t | 6/16 |
| -r | -t | 6/16 |

Causality?

- When Bayes' nets reflect the true causal patterns:
 - Often simpler (nodes have fewer parents)
 - Often easier to think about
 - Often easier to elicit from experts
- BNs need not actually be causal
 - Sometimes no causal net exists over the domain (especially if variables are missing)
 - E.g. consider the variables *Traffic* and *Drips*
 - End up with arrows that reflect correlation, not causation
- What do the arrows really mean?
 - Topology may happen to encode causal structure
 - **Topology really encodes conditional independence**



$$P(x_i | x_1, \dots, x_{i-1}) = P(x_i | \text{parents}(X_i))$$

Bayes' Nets

- So far: how a Bayes' net encodes a joint distribution
- Next: how to answer queries about that distribution
 - Today:
 - First assembled BNs using an intuitive notion of conditional independence as causality
 - Then saw that key property is conditional independence
 - Main goal: answer queries about conditional independence and influence
- After that: how to answer numerical queries (inference)

