


# Announcements: Midterm

---

- **This Thursday, October 17, 2024, 7-9 PM PT.**
  - If you need to take the alternate exam (same day, 9-11 PM PT, in-person only) or if you need DSP accommodations and you have not filled out the alternate exam request form, send us an email at [cs188@berkeley.edu](mailto:cs188@berkeley.edu) immediately.
- **Topics:**
  - Search, CSPs, Games, MDPs, Reinforcement Learning, and Bayes Nets (only up to representation). These were covered in: Lectures 1-13, Notes 1-6.4, Discussions 1-5, Projects 1-3, and Homeworks 1-5.
- **Format:**
  - Closed-book, closed-notes, and closed-internet. No calculators are allowed (no questions require a calculator). However, you may use 1 cheat sheet (two-sided) of your own design, handwritten or typed.
- **Room Assignment: was emailed to you by Saturday, October 12th**
  - If you did not receive an email yet, or if you are supposed to be taking an alternate exam and your email did not indicate that, send an email to [cs188@berkeley.edu](mailto:cs188@berkeley.edu).
  - Please try to arrive early, so that we can start on time.
- **Check Ed in case of any further announcements**

# CS 188: Artificial Intelligence

## Bayes' Nets: Independence



Instructors: Pieter Abbeel & Igor Mordatch --- University of California, Berkeley

[Many of these slides were originally created by Dan Klein and Pieter Abbeel for CS188 Intro to AI at UC Berkeley.]

# Probability Recap


- Conditional probability  $P(x|y) = \frac{P(x, y)}{P(y)}$
- Product rule  $P(x, y) = P(x|y)P(y)$
- Chain rule 
$$\begin{aligned} P(X_1, X_2, \dots, X_n) &= P(X_1)P(X_2|X_1)P(X_3|X_1, X_2) \dots \\ &= \prod_{i=1}^n P(X_i|X_1, \dots, X_{i-1}) \end{aligned}$$
- X, Y independent if and only if:  $\forall x, y : P(x, y) = P(x)P(y)$
- X and Y are conditionally independent given Z if and only if:

$$\forall x, y, z : P(x, y|z) = P(x|z)P(y|z)$$

$$X \perp\!\!\!\perp Y | Z$$

# Bayes' Nets

- A Bayes' net is an efficient encoding of a probabilistic model of a domain



- Questions we can ask:

- Modeling: what BN is most appropriate for a given domain?
- Representation: given a BN graph, what kinds of distributions can it encode?
- Inference: given a fixed BN, what is  $P(X \mid e)$ ?



# Bayes' Net Semantics

- A directed, acyclic graph, one node per random variable
- A conditional probability table (CPT) for each node
  - A collection of distributions over  $X$ , one for each combination of parents' values

$$P(X|a_1 \dots a_n)$$


- Bayes' nets implicitly encode joint distributions
  - As a product of local conditional distributions
  - To see what probability a BN gives to a full assignment, multiply all the relevant conditionals together:

$$P(x_1, x_2, \dots, x_n) = \prod_{i=1}^n P(x_i | \text{parents}(X_i))$$




# Example: Alarm Network

B	P(B)
+b	0.001
-b	0.999



E	P(E)
+e	0.002
-e	0.998

A	M	P(M A)
+a	+m	0.7
+a	-m	0.3
-a	+m	0.01
-a	-m	0.99




A	J	P(J A)
+a	+j	0.9
+a	-j	0.1
-a	+j	0.05
-a	-j	0.95

B	E	A	P(A B,E)
+b	+e	+a	0.95
+b	+e	-a	0.05
+b	-e	+a	0.94
+b	-e	-a	0.06
-b	+e	+a	0.29
-b	+e	-a	0.71
-b	-e	+a	0.001
-b	-e	-a	0.999

$$P(+b, -e, +a, -j, +m) =$$


# Example: Alarm Network

B	P(B)
+b	0.001
-b	0.999



E	P(E)
+e	0.002
-e	0.998

A	M	P(M A)
+a	+m	0.7
+a	-m	0.3
-a	+m	0.01
-a	-m	0.99



A	J	P(J A)
+a	+j	0.9
+a	-j	0.1
-a	+j	0.05
-a	-j	0.95

B	E	A	P(A B,E)
+b	+e	+a	0.95
+b	+e	-a	0.05
+b	-e	+a	0.94
+b	-e	-a	0.06
-b	+e	+a	0.29
-b	+e	-a	0.71
-b	-e	+a	0.001
-b	-e	-a	0.999

$$\begin{aligned}
 P(+b, -e, +a, -j, +m) &= \\
 P(+b)P(-e)P(+a|+b, -e)P(-j|+a)P(+m|+a) &= \\
 0.001 \times 0.998 \times 0.94 \times 0.1 \times 0.7 &
 \end{aligned}$$

# Size of a Bayes' Net

- How big is a joint distribution over  $N$  Boolean variables?

$$2^N$$



- How big is an  $N$ -node net if nodes have up to  $k$  parents?

$$O(N * 2^{k+1})$$

- Both give you the power to calculate

$$P(X_1, X_2, \dots, X_n)$$

- BNs: Huge space savings!
- Also easier to elicit local CPTs
- Also faster to answer queries (coming)





# Bayes' Nets

---

- ✓ Representation
  - Conditional Independences
  - Probabilistic Inference
  - Learning Bayes' Nets from Data

# Conditional Independence

- X and Y are **independent** if

$$\forall x, y \quad P(x, y) = P(x)P(y) \quad \text{---} \rightarrow \quad X \perp\!\!\!\perp Y$$


- X and Y are **conditionally independent** given Z

$$\forall x, y, z \quad P(x, y|z) = P(x|z)P(y|z) \quad \text{---} \rightarrow \quad X \perp\!\!\!\perp Y|Z$$

- (Conditional) independence is a property of a distribution

- Example:

$$\textit{Alarm} \perp\!\!\!\perp \textit{Fire} | \textit{Smoke}$$




# Bayes Nets: Assumptions

- Assumptions we are required to make to define the Bayes net when given the graph:


$$P(x_i | x_1 \cdots x_{i-1}) = P(x_i | \text{parents}(X_i))$$

- Beyond above “chain rule  $\rightarrow$  Bayes net” conditional independence assumptions
  - Often additional conditional independences
  - They can be read off the graph
- Important for modeling: understand assumptions made when choosing a Bayes net graph



# Example

---




- Conditional independence assumptions directly from simplifications in chain rule:
- Additional implied conditional independence assumptions?

# Independence in a BN

- Important question about a BN:
  - Are two nodes independent given certain evidence?
    - If yes, can prove using algebra (tedious in general)
    - If no, can prove with a counter example


- Example:



- Question: for this BN graph, are X and Z necessarily independent?\*
- Answer: no. Example: low pressure causes rain, which causes traffic.
- X can influence Z, Z can influence X (via Y)
- Addendum: they *could* be independent: how?

\* Equivalent phrasing: for this BN graph, are X and Z guaranteed to be independent no matter the choice of CPTs

# D-separation



# D-separation: Overview

---

- **D-separation:**
  - a condition / algorithm for answering conditional independence queries from just studying the graph
- **How:**
  - Study independence properties for triples
  - Analyze complex cases as composition of triples

# Triple Type 1: Causal Chains

- This configuration is a “causal chain”



X: Low pressure

Y: Rain

Z: Traffic

$$P(x, y, z) = P(x)P(y|x)P(z|y)$$

- Guaranteed X independent of Z ? *No!*

- One example set of CPTs for which X is not independent of Z is sufficient to show this independence is not guaranteed.

- Example:

- Low pressure causes rain causes traffic, high pressure causes no rain causes no traffic

- In numbers:

$$P(+y \mid +x) = 1, P(-y \mid -x) = 1, \\ P(+z \mid +y) = 1, P(-z \mid -y) = 1$$



# Triple Type 1: Causal Chains

- This configuration is a “causal chain”

- Guaranteed X independent of Z given Y?



X: Low pressure

Y: Rain

Z: Traffic

$$P(x, y, z) = P(x)P(y|x)P(z|y)$$


$$\begin{aligned} P(z|x, y) &= \frac{P(x, y, z)}{P(x, y)} \\ &= \frac{P(x)P(y|x)P(z|y)}{P(x)P(y|x)} \\ &= P(z|y) \end{aligned}$$

*Yes!*

- Evidence along the chain “blocks” the influence

# Triple Type 2: Common Cause

- This configuration is a “common cause”



$$P(x, y, z) = P(y)P(x|y)P(z|y)$$

- Guaranteed X independent of Z? **No!**

- One example set of CPTs for which X is not independent of Z is sufficient to show this independence is not guaranteed.

- Example:

- Project due causes both forums busy and lab full


- In numbers:

$$P(+x | +y) = 1, P(-x | -y) = 1, \\ P(+z | +y) = 1, P(-z | -y) = 1$$

# Triple Type 2: Common Cause

- This configuration is a “common cause”

- Guaranteed X and Z independent given Y?



$$P(x, y, z) = P(y)P(x|y)P(z|y)$$


$$\begin{aligned} P(z|x, y) &= \frac{P(x, y, z)}{P(x, y)} \\ &= \frac{P(y)P(x|y)P(z|y)}{P(y)P(x|y)} \\ &= P(z|y) \end{aligned}$$

*Yes!*

- Observing the cause blocks influence between effects.

# Triple Type 3: Common Effect

- Last configuration: two causes of one effect (v-structures)




- Are X and Y independent?
  - **Yes**: the ballgame and the rain cause traffic, but they are not correlated
  - Still need to prove they must be (try it!)
- Are X and Y independent given Z?
  - **No**: seeing traffic puts the rain and the ballgame in competition as explanation.
- **This is backwards from the other cases**
  - Observing an effect **activates** influence between possible causes.

# Recap of Triples



## Active Triples

## Inactive Triples




Causal Chain:



Common Cause:



Common Effect ("v-structure")




# The General Case

---



# The General Case

- General question: in a given BN, are two variables independent (given evidence)?
- Solution: analyze the graph
- Each path can be seen as repetitions of the three canonical cases



# From Triples to Paths to D-Separation

- A path is active if each (overlapping) triple is active:

Note: e.g. for a path  $A - B - C - D - E$ , the triples are:

$A - B - C$ ,  $B - C - D$ ,  $C - D - E$

Note: all it takes to block a path is a single inactive segment

- Are  $X$  and  $Y$  “D-separated” given evidence variables  $\{Z\}$ ?

- Consider all (undirected) paths from  $X$  to  $Y$
- If none of the paths are active, then  $X$  and  $Y$  are D-separated given  $\{Z\}$
- On the other hand, if there is at least one active path, then  $X$  and  $Y$  are not D-separated given  $\{Z\}$

- Independence and D-separation:





$X$  and  $Y$  are guaranteed conditionally independent given  $\{Z\}$

IF AND ONLY IF



$X$  and  $Y$  are d-separated given  $\{Z\}$

→ just need to check the graph

Active Triples



Inactive Triples






# Example

---

$R \perp\!\!\!\perp B$       *Yes*

$R \perp\!\!\!\perp B | T$

$R \perp\!\!\!\perp B | T'$



# Example


$L \perp\!\!\!\perp T' \mid T$     *Yes*

$L \perp\!\!\!\perp B$     *Yes*

$L \perp\!\!\!\perp B \mid T$

$L \perp\!\!\!\perp B \mid T'$

$L \perp\!\!\!\perp B \mid T, R$     *Yes*




# Example

- Variables:
  - R: Raining
  - T: Traffic
  - D: Roof drips
  - S: I'm sad
- Questions:

$$T \perp\!\!\!\perp D$$

$$T \perp\!\!\!\perp D \mid R \quad \text{Yes}$$

$$T \perp\!\!\!\perp D \mid R, S$$




# Structure Implications

- Given a Bayes net structure, can run d-separation algorithm to build a complete list of conditional independences that are necessarily true of the form



$$X_i \perp\!\!\!\perp X_j \mid \{X_{k_1}, \dots, X_{k_n}\}$$

- This list determines the set of probability distributions that can be represented



# Computing All Independences





COMPUTE ALL THE  
INDEPENDENCES!



# Topology Limits Distributions

- Given some graph topology  $G$ , only certain joint distributions can be encoded
- The graph structure guarantees certain (conditional) independences
- (There might be more independence)
- Adding arcs increases the set of distributions, but has several costs
- Full conditioning can encode any distribution

$$\{X \perp\!\!\!\perp Y, X \perp\!\!\!\perp Z, Y \perp\!\!\!\perp Z, \\ X \perp\!\!\!\perp Z \mid Y, X \perp\!\!\!\perp Y \mid Z, Y \perp\!\!\!\perp Z \mid X\}$$



# Bayes Nets Representation Summary

---

- Bayes nets compactly encode joint distributions
- Guaranteed independencies of distributions can be deduced from BN graph structure
- D-separation gives precise conditional independence guarantees from graph alone
- A Bayes' net's joint distribution may have further (conditional) independence that is not detectable until you inspect its specific distribution

# Bayes' Nets

---

✓ Representation

✓ Conditional Independences

- Probabilistic Inference

- Enumeration (exact, exponential complexity)

- Variable elimination (exact, worst-case exponential complexity, often better)

- Probabilistic inference is NP-complete

- Sampling (approximate)

- Learning Bayes' Nets from Data