

# CS 188: Artificial Intelligence Spring 2010

## Lecture 15: Bayes' Nets II – Independence 3/9/2010

Pieter Abbeel – UC Berkeley

Many slides over the course adapted from Dan Klein, Stuart Russell,  
Andrew Moore

## Announcements

---

- **Current readings**
  - Require login
- **Assignments**
  - W4 due Thursday
- **Midterm**
  - 3/18, 6-9pm, 0010 Evans --- no lecture on 3/18
  - We will be posting practice midterms
  - One page note sheet, non-programmable calculators
  - Topics go through Thursday, not next Tuesday

## Outline

---

- Thus far: Probability
- Today: Bayes nets
  - Semantics
  - (Conditional) Independence

3

## Probability recap

---

- Conditional probability  $P(x|y) = \frac{P(x,y)}{P(y)}$
- Product rule  $P(x,y) = P(x|y)P(y)$
- Chain rule  $P(X_1, X_2, \dots, X_n) = P(X_1)P(X_2|X_1)P(X_3|X_1, X_2) \dots$
- X, Y independent iff:  $\forall x, y : P(x, y) = P(x)P(y)$
- X and Y are conditionally independent given Z iff:  
 $\forall x, y, z : P(x, y|z) = P(x|z)P(y|z)$   $X \perp\!\!\!\perp Y | Z$  4


# Bayes' Nets: Big Picture

- Two problems with using full joint distribution tables as our probabilistic models:
  - Unless there are only a few variables, the joint is WAY too big to represent explicitly
  - Hard to learn (estimate) anything empirically about more than a few variables at a time
- Bayes' nets:** a technique for describing complex joint distributions (models) using simple, local distributions (conditional probabilities)
  - More properly called **graphical models**
  - We describe how variables locally interact
  - Local interactions chain together to give global, indirect interactions
  - For about 10 min, we'll be vague about how these interactions are specified

5

# Graphical Model Notation

- Nodes: variables (with domains)**
  - Can be assigned (observed) or unassigned (unobserved)
- Arcs: interactions**
  - Similar to CSP constraints
  - Indicate "direct influence" between variables
  - Formally: encode conditional independence (more later)
- For now: imagine that arrows mean direct causation (in general, they don't!)**



8

## Example: Coin Flips

---

- N independent coin flips




- No interactions between variables:  
**absolute independence**

9

## Example: Traffic

---

- Variables:
  - R: It rains
  - T: There is traffic
- Model 1: independence
- Model 2: rain causes traffic
- Why is an agent using model 2 better?



10

## Example: Traffic II

---

- Let's build a causal graphical model
- Variables
  - T: Traffic
  - R: It rains
  - L: Low pressure
  - D: Roof drips
  - B: Ballgame
  - C: Cavity

11

## Example: Alarm Network


---

- Variables
  - B: Burglary
  - A: Alarm goes off
  - M: Mary calls
  - J: John calls
  - E: Earthquake!

12

# Bayes' Net Semantics

- Let's formalize the semantics of a Bayes' net
- A set of nodes, one per variable  $X$
- A directed, acyclic graph
- A conditional distribution for each node
  - A collection of distributions over  $X$ , one for each combination of parents' values



$$P(X|A_1 \dots A_n)$$


$$P(X|a_1 \dots a_n)$$

- CPT: conditional probability table
- Description of a noisy "causal" process

*A Bayes net = Topology (graph) + Local Conditional Probabilities*

13

# Probabilities in BNs



- Bayes' nets **implicitly** encode joint distributions
  - As a product of local conditional distributions
  - To see what probability a BN gives to a full assignment, multiply all the relevant conditionals together:

$$P(x_1, x_2, \dots, x_n) = \prod_{i=1}^n P(x_i | \text{parents}(X_i))$$


- Example:

$$P(+cavity, +catch, -toothache)$$

- This lets us reconstruct any entry of the full joint
- Not every BN can represent every joint distribution
  - The topology enforces certain conditional independencies

14

## Example: Coin Flips




$$P(h, h, t, h) =$$

*Only distributions whose variables are absolutely independent can be represented by a Bayes' net with no arcs.*

15

## Example: Traffic



16

## Example: Alarm Network

B	P(B)
+b	0.001
-b	0.999



E	P(E)
+e	0.002
-e	0.998

A	J	P(J A)
+a	+j	0.9
+a	-j	0.1
-a	+j	0.05
-a	-j	0.95

A	M	P(M A)
+a	+m	0.7
+a	-m	0.3
-a	+m	0.01
-a	-m	0.99

B	E	A	P(A B,E)
+b	+e	+a	0.95
+b	+e	-a	0.05
+b	-e	+a	0.94
+b	-e	-a	0.06
-b	+e	+a	0.29
-b	+e	-a	0.71
-b	-e	+a	0.001
-b	-e	-a	0.999

## Size of a Bayes' Net

- How big is a joint distribution over  $N$  Boolean variables?  
 $2^N$
- How big is an  $N$ -node net if nodes have up to  $k$  parents?  
 $O(N * 2^{k+1})$
- Both give you the power to calculate  $P(X_1, X_2, \dots, X_n)$
- BNs: Huge space savings!
- Also easier to elicit local CPTs
- Also turns out to be faster to answer queries (coming)



# Bayes' Nets

---

- So far: how a Bayes' net encodes a joint distribution
- Next: how to answer queries about that distribution
  - Key idea: conditional independence
- After that: how to answer numerical queries (inference) more efficiently than by first constructing the joint distribution

19

# Conditional Independence


---

- **Reminder: independence**
  - X and Y are **independent** if
$$\forall x, y \quad P(x, y) = P(x)P(y) \quad \dashrightarrow \quad X \perp\!\!\!\perp Y$$
  - X and Y are **conditionally independent** given Z
$$\forall x, y, z \quad P(x, y|z) = P(x|z)P(y|z) \quad \dashrightarrow \quad X \perp\!\!\!\perp Y | Z$$
- (Conditional) independence is a property of a distribution

23


## Example: Independence

- For this graph, you can fiddle with  $\theta$  (the CPTs) all you want, but you won't be able to represent any distribution in which the flips are dependent!




## Topology Limits Distributions

- Given some graph topology  $G$ , only certain joint distributions can be encoded
- The graph structure guarantees certain (conditional) independences
- (There might be more independence)
- Adding arcs increases the set of distributions, but has several costs
- Full conditioning can encode any distribution



## Independence in a BN


- Important question about a BN:
  - Are two nodes independent given certain evidence?
  - If yes, can prove using algebra (tedious in general)
  - If no, can prove with a counter example
  - Example:



- Question: are X and Z necessarily independent?
  - Answer: no. Example: low pressure causes rain, which causes traffic.
  - X can influence Z, Z can influence X (via Y)
  - Addendum: they *could* be independent: how?

## Causal Chains

- This configuration is a “causal chain”



X: Low pressure

Y: Rain

Z: Traffic

$$P(x, y, z) = P(x)P(y|x)P(z|y)$$

- Is X independent of Z given Y?

$$\begin{aligned} P(z|x, y) &= \frac{P(x, y, z)}{P(x, y)} = \frac{P(x)P(y|x)P(z|y)}{P(x)P(y|x)} \\ &= P(z|y) \quad \text{Yes!} \end{aligned}$$

- Evidence along the chain “blocks” the influence


27

## Common Cause

- Another basic configuration: two effects of the same cause

- Are X and Z independent?
- Are X and Z independent given Y?

$$P(z|x, y) = \frac{P(x, y, z)}{P(x, y)} = \frac{P(y)P(x|y)P(z|y)}{P(y)P(x|y)} = P(z|y) \quad \text{Yes!}$$



Y: Project due  
X: Newsgroup busy  
Z: Lab full


- Observing the cause blocks influence between effects.

28

## Common Effect

- Last configuration: two causes of one effect (v-structures)

- Are X and Z independent?
  - Yes: the ballgame and the rain cause traffic, but they are not correlated
  - Still need to prove they must be (try it!)
- Are X and Z independent given Y?
  - No: seeing traffic puts the rain and the ballgame in competition as explanation?
- This is backwards from the other cases
  - Observing an effect **activates** influence between possible causes.



X: Raining  
Z: Ballgame  
Y: Traffic

29

## The General Case

---

- Any complex example can be analyzed using these three canonical cases
- General question: in a given BN, are two variables independent (given evidence)?
- Solution: analyze the graph

30