

CS 188: Artificial Intelligence Spring 2010

Lecture 17: Bayes' Nets IV – Inference 3/16/2010

Pieter Abbeel – UC Berkeley

Many slides over this course adapted from Dan Klein, Stuart Russell,
Andrew Moore

Announcements

- **Assignments**
 - W4 back today in lecture
 - Any assignments you have not picked up yet
 - In bin in 283 Soda [same room as for submission drop-off]

- **Midterm**
 - 3/18, 6-9pm, 0010 Evans --- no lecture on Thursday
 - We have posted practice midterms (and finals)
 - One note letter-size note sheet (two sides), non-programmable calculators [strongly encouraged to compose your own!]
 - Topics go through last Thursday
 - **Section this week: midterm review**

2


Bayes' Net Semantics

- Let's formalize the semantics of a Bayes' net
- A set of nodes, one per variable X
- A directed, acyclic graph
- A conditional distribution for each node
 - A collection of distributions over X , one for each combination of parents' values

$$P(X|a_1 \dots a_n)$$

- CPT: conditional probability table
- Description of a noisy "causal" process

A Bayes net = Topology (graph) + Local Conditional Probabilities



3

Probabilities in BNs

- For all joint distributions, we have (chain rule):

$$P(x_1, x_2, \dots, x_n) = \prod_{i=1}^n P(x_i | x_1, \dots, x_{i-1})$$

- Bayes' nets **implicitly** encode joint distributions
 - As a product of local conditional distributions
 - To see what probability a BN gives to a full assignment, multiply all the relevant conditionals together:

$$P(x_1, x_2, \dots, x_n) = \prod_{i=1}^n P(x_i | \text{parents}(X_i))$$

- This lets us reconstruct any entry of the full joint
- Not every BN can represent every joint distribution
 - The topology enforces certain conditional independencies

4

Bayes Nets Representation Summary

- Bayes nets compactly encode joint distributions
- Guaranteed independencies of distributions can be deduced from BN graph structure
- D-separation gives precise conditional independence guarantees from graph alone
- A Bayes' net's joint distribution may have further (conditional) independence that is not detectable until you inspect its specific distribution

5


Inference

- Inference: calculating some useful quantity from a joint probability distribution
- Examples:
 - Posterior probability:

$$P(Q|E_1 = e_1, \dots, E_k = e_k)$$

- Most likely explanation:

$$\operatorname{argmax}_q P(Q = q|E_1 = e_1 \dots)$$




6

Inference by Enumeration

- Given unlimited time, inference in BNs is easy
- Recipe:
 - State the marginal probabilities you need
 - Figure out ALL the atomic probabilities you need
 - Calculate and combine them
- Example:

$$P(+b | +j, +m) =$$

$$\frac{P(+b, +j, +m)}{P(+j, +m)}$$



7

Example: Enumeration


- In this simple method, we only need the BN to synthesize the joint entries

$$P(+b, +j, +m) =$$

$$\begin{aligned}
 &P(+b)P(+e)P(+a|+b, +e)P(+j|+a)P(+m|+a)+ \\
 &P(+b)P(+e)P(-a|+b, +e)P(+j|-a)P(+m|-a)+ \\
 &P(+b)P(-e)P(+a|+b, -e)P(+j|+a)P(+m|+a)+ \\
 &P(+b)P(-e)P(-a|+b, -e)P(+j|-a)P(+m|-a)
 \end{aligned}$$

8

Inference by Enumeration?



9

Variable Elimination

- Why is inference by enumeration so slow?
 - You join up the whole joint distribution before you sum out the hidden variables
 - You end up repeating a lot of work!
- Idea: interleave joining and marginalizing!
 - Called “Variable Elimination”
 - Still NP-hard, but usually much faster than inference by enumeration
- We’ll need some new notation to define VE

10

Factor Zoo I

- Joint distribution: $P(X,Y)$

- Entries $P(x,y)$ for all x, y
- Sums to 1

$P(T, W)$

T	W	P
hot	sun	0.4
hot	rain	0.1
cold	sun	0.2
cold	rain	0.3

- Selected joint: $P(x,Y)$

- A slice of the joint distribution
- Entries $P(x,y)$ for fixed x , all y
- Sums to $P(x)$

$P(cold, W)$

T	W	P
cold	sun	0.2
cold	rain	0.3

11

Factor Zoo II

- Family of conditionals:

$P(X|Y)$

- Multiple conditionals
- Entries $P(x|y)$ for all x, y
- Sums to $|Y|$

$P(W|T)$

T	W	P
hot	sun	0.8
hot	rain	0.2
cold	sun	0.4
cold	rain	0.6

$P(W|hot)$

$P(W|cold)$

- Single conditional: $P(Y|x)$

- Entries $P(y|x)$ for fixed x , all y
- Sums to 1

$P(W|cold)$

T	W	P
cold	sun	0.4
cold	rain	0.6

12

Factor Zoo III

- Specified family: $P(y | X)$
 - Entries $P(y | x)$ for fixed y , but for all x
 - Sums to ... who knows!

$P(\text{rain}|T)$

T	W	P
hot	rain	0.2
cold	rain	0.6


} $P(\text{rain}|\text{hot})$
} $P(\text{rain}|\text{cold})$

- In general, when we write $P(Y_1 \dots Y_N | X_1 \dots X_M)$
 - It is a “factor,” a multi-dimensional array
 - Its values are all $P(y_1 \dots y_N | x_1 \dots x_M)$
 - Any assigned X or Y is a dimension missing (selected) from the array

13

Example: Traffic Domain

- Random Variables
 - R: Raining
 - T: Traffic
 - L: Late for class!



$P(R)$

+r	0.1
-r	0.9

$P(T|R)$

+r	+t	0.8
+r	-t	0.2
-r	+t	0.1
-r	-t	0.9

$P(L|R)$

+t	+l	0.3
+t	-l	0.7
-t	+l	0.1
-t	-l	0.9

14

Variable Elimination Outline

- Track objects called **factors**
- Initial factors are local CPTs (one per node)

+r	0.1
-r	0.9

+r	+t	0.8
+r	-t	0.2
-r	+t	0.1
-r	-t	0.9

+t	+l	0.3
+t	-l	0.7
-t	+l	0.1
-t	-l	0.9

- Any known values are selected
 - E.g. if we know $L = +l$, the initial factors are

+r	0.1
-r	0.9

+r	+t	0.8
+r	-t	0.2
-r	+t	0.1
-r	-t	0.9


+t	+l	0.3
-t	+l	0.1

- VE: Alternately join factors and eliminate variables

15

Operation 1: Join Factors

- First basic operation: **joining factors**
- Combining factors:
 - Just like a database join
 - Get all factors over the joining variable
 - Build a new factor over the union of the variables involved
- Example: Join on R




- Computation for each entry: pointwise products

$$\forall r, t : P(r, t) = P(r) \cdot P(t|r)$$


16

Example: Multiple Joins



18

Example: Multiple Joins



19

Operation 2: Eliminate

- Second basic operation: **marginalization**
- Take a factor and sum out a variable
 - Shrinks a factor to a smaller one
 - A **projection** operation
- Example:

$P(R, T)$			sum R	$P(T)$	
+r	+t	0.08	→	+t	0.17
+r	-t	0.02		-t	0.83
-r	+t	0.09			
-r	-t	0.81			


20

Multiple Elimination


R, T, L		T, L		L								
$P(R, T, L)$												
+r	+t	+l	0.024	Sum out R	→	$P(T, L)$			Sum out T	→	$P(L)$	
+r	+t	-l	0.056	+t		+l	0.051	+l	0.134			
+r	-t	+l	0.002	+t	-l	0.119	-l	0.886				
+r	-t	-l	0.018	-t	+l	0.083						
-r	+t	+l	0.027	-t	-l	0.747						
-r	+t	-l	0.063									
-r	-t	+l	0.081									
-r	-t	-l	0.729									

21

P(L) : Marginalizing Early!



Marginalizing Early (aka VE*)



Evidence

- If evidence, start with factors that select that evidence

- No evidence uses these initial factors:

+r	0.1
-r	0.9

+r	+t	0.8
+r	-t	0.2
-r	+t	0.1
-r	-t	0.9

+t	+l	0.3
+t	-l	0.7
-t	+l	0.1
-t	-l	0.9

- Computing $P(L|+r)$, the initial factors become:

+r	0.1
----	-----

+r	+t	0.8
+r	-t	0.2

+t	+l	0.3
+t	-l	0.7
-t	+l	0.1
-t	-l	0.9

- We eliminate all vars other than query + evidence

24


Evidence II

- Result will be a selected joint of query and evidence

- E.g. for $P(L|+r)$, we'd end up with:

+r	+l	0.026
+r	-l	0.074

Normalize



+l	0.26
-l	0.74

- To get our answer, just normalize this!

- That's it!

25


General Variable Elimination

- Query: $P(Q|E_1 = e_1, \dots, E_k = e_k)$
- Start with initial factors:
 - Local CPTs (but instantiated by evidence)
- While there are still hidden variables (not Q or evidence):
 - Pick a hidden variable H
 - Join all factors mentioning H
 - Eliminate (sum out) H
- Join all remaining factors and normalize

26

Variable Elimination Bayes Rule


Start / Select



$P(A|B) \rightarrow P(a|B)$

B	A	P
+b	+a	0.8
b	-a	0.2
-b	+a	0.1
-b	-a	0.9

Join on B



$P(a, B)$

A	B	P
+a	+b	0.08
+a	-b	0.09

Normalize

$P(B|a)$

A	B	P
+a	+b	8/17
+a	-b	9/17

27

Example

$$P(B|j, m) \propto P(B, j, m)$$

$P(B)$	$P(E)$	$P(A B, E)$	$P(j A)$	$P(m A)$
--------	--------	-------------	----------	----------

Choose A

$$\begin{array}{l}
 P(A|B, E) \\
 P(j|A) \\
 P(m|A)
 \end{array}
 \xrightarrow{\times}
 P(j, m, A|B, E)
 \xrightarrow{\Sigma}
 P(j, m|B, E)$$

$P(B)$	$P(E)$	$P(j, m B, E)$
--------	--------	----------------

28

Example

$P(B)$	$P(E)$	$P(j, m B, E)$
--------	--------	----------------

Choose E

$$\begin{array}{l}
 P(E) \\
 P(j, m|B, E)
 \end{array}
 \xrightarrow{\times}
 P(j, m, E|B)
 \xrightarrow{\Sigma}
 P(j, m|B)$$

$P(B)$	$P(j, m B)$
--------	-------------

Finish with B

$$\begin{array}{l}
 P(B) \\
 P(j, m|B)
 \end{array}
 \xrightarrow{\times}
 P(j, m, B)
 \xrightarrow{\text{Normalize}}
 P(B|j, m)$$

29

Variable Elimination

- What you need to know:
 - Should be able to run it on small examples, understand the factor creation / reduction flow
 - Better than enumeration: saves time by marginalizing variables as soon as possible rather than at the end
- We will see special cases of VE later
 - On tree-structured graphs, variable elimination runs in polynomial time, like tree-structured CSPs
 - You'll have to implement a tree-structured special case to track invisible ghosts (Project 4)