

CS 188: Artificial Intelligence

Hidden Markov Models



University of California, Berkeley


Announcements

- HW6 self-assessment
 - Due on Friday 3/21/25 at 11:59 PT
- Enjoy Spring Break!

Reasoning over Time or Space

- Often, we want to reason about a sequence of observations
 - Speech recognition
 - Robot localization
 - User attention
 - Medical monitoring
 - Language understanding
- Need to introduce time (or space) into our models and update beliefs based on:
 - Getting more evidence (we did this with BNs)
 - World changing over time (new this week)

Motivating Example: Pacman Sonar



Today's Topics

- Quick probability recap
- **Markov Chains & their Stationary Distributions**
 - How beliefs about state change with passage of time
- **Hidden Markov Models (HMMs) formulation**
 - How beliefs change with passage of time and evidence
- **Filtering with HMMs**
 - How to infer beliefs from evidence

Probability Recap

- Conditional probability

$$P(x|y) = \frac{P(x, y)}{P(y)}$$

- Marginal probability

$$P(x) = \sum_y P(x, y)$$

- Product rule

$$P(x, y) = P(x|y)P(y)$$

- Chain rule

$$\begin{aligned} P(X_1, X_2, \dots, X_n) &= P(X_1)P(X_2|X_1)P(X_3|X_1, X_2) \dots \\ &= \prod_{i=1}^n P(X_i|X_1, \dots, X_{i-1}) \end{aligned}$$

Probability Recap


- X, Y independent if and only if: $\forall x, y : P(x, y) = P(x)P(y)$
- X and Y are conditionally independent given Z : $X \perp\!\!\!\perp Y | Z$ if and only if:
$$\forall x, y, z : P(x, y|z) = P(x|z)P(y|z)$$
- Proportionality: $P(X) \propto f(X)$ or $P(X) \propto_x f(X)$ means $P(X) = kf(X)$ (for some constant k that doesn't depend on X). Equivalent to: $P(X) = \frac{f(X)}{\sum_x f(x)}$

■ Example:

| X | $\propto f(X)$ | $P(X)$ |
|-------|----------------|---------------------|
| x_1 | 0.4 | $0.4 / (0.4 + 0.2)$ |
| x_2 | 0.2 | $0.2 / (0.4 + 0.2)$ |

Markov Models


- Value of X at a given time is called the **state**



$$P(X_1) \quad P(X_t|X_{t-1})$$

- Parameters: called **transition probabilities** or dynamics, specify how the state evolves over time (also, initial state probabilities)
- Stationarity assumption: transition probabilities the same at all times
- Same as MDP transition model, but no choice of action
- A "growable" BN (can always use BN methods if we truncate to fixed length)


Conditional Independence



- **Basic conditional independence:**
 - Past and future independent given the present
 - Each time step only depends on the previous
 - This is called the (first order) Markov property



Example Markov Chain: Weather

- States: $X = \{\text{rain, sun}\}$
- Initial distribution: 1.0 sun
- CPT $P(X_t | X_{t-1})$:



Two new ways of representing the same CPT

| X_{t-1} | X_t | $P(X_t X_{t-1})$ |
|-----------|-------|--------------------|
| sun | sun | 0.9 |
| sun | rain | 0.1 |
| rain | sun | 0.3 |
| rain | rain | 0.7 |



Example Markov Chain: Weather

- Initial distribution: 1.0 sun

- We know: $P(X_1)$ $P(X_t|X_{t-1})$

| X_{t-1} | X_t | $P(X_t X_{t-1})$ |
|-----------|-------|------------------|
| sun | sun | 0.9 |
| sun | rain | 0.1 |
| rain | sun | 0.3 |
| rain | rain | 0.7 |

- What is the probability distribution after one step?

$$\begin{aligned} P(X_2 = sun) &= \sum_{x_1} P(x_1, X_2 = sun) = \sum_{x_1} P(X_2 = sun|x_1)P(x_1) \\ &= P(X_2 = sun|X_1 = sun)P(X_1 = sun) + \\ &\quad P(X_2 = sun|X_1 = rain)P(X_1 = rain) \\ &= 0.9 \cdot 1.0 + 0.3 \cdot 0.0 = 0.9 \end{aligned}$$

Example Markov Chain: Weather

- Initial distribution: 1.0 sun
 - We know: $P(X_1)$ $P(X_t|X_{t-1})$

| X_{t-1} | X_t | $P(X_t X_{t-1})$ |
|-----------|-------|------------------|
| sun | sun | 0.9 |
| sun | rain | 0.1 |
| rain | sun | 0.3 |
| rain | rain | 0.7 |


- In matrix form:

$$\text{row} = X_t, \text{ col} = X_{t-1}$$
$$P(X_2) = \begin{bmatrix} 0.9 & 0.3 \\ 0.1 & 0.7 \end{bmatrix} P(X_1)$$

$$\begin{bmatrix} 0.9 \\ 0.1 \end{bmatrix} = \begin{bmatrix} 0.9 & 0.3 \\ 0.1 & 0.7 \end{bmatrix} \begin{bmatrix} 1.0 \\ 0 \end{bmatrix}$$

Mini-Forward Algorithm

- Question: What's $P(X)$ on some day t ?




- We know $P(X_1)$ and $P(X_t|X_{t-1})$

$$P(X_1) = \text{known}$$

$$\begin{aligned} P(x_t) &= \sum_{x_{t-1}} P(x_{t-1}, x_t) \\ &= \sum_{x_{t-1}} P(x_t | x_{t-1}) P(x_{t-1}) \end{aligned}$$

Forward simulation



Example Run of Mini-Forward Algorithm

- From initial observation of sun

$$\begin{array}{ccccccc} \left\langle \begin{array}{c} 1.0 \\ 0.0 \end{array} \right\rangle & \left\langle \begin{array}{c} 0.9 \\ 0.1 \end{array} \right\rangle & \left\langle \begin{array}{c} 0.84 \\ 0.16 \end{array} \right\rangle & \left\langle \begin{array}{c} 0.804 \\ 0.196 \end{array} \right\rangle & \longrightarrow & \left\langle \begin{array}{c} 0.75 \\ 0.25 \end{array} \right\rangle \\ P(X_1) & P(X_2) & P(X_3) & P(X_4) & & P(X_\infty) \end{array}$$


- From initial observation of rain

$$\begin{array}{ccccccc} \left\langle \begin{array}{c} 0.0 \\ 1.0 \end{array} \right\rangle & \left\langle \begin{array}{c} 0.3 \\ 0.7 \end{array} \right\rangle & \left\langle \begin{array}{c} 0.48 \\ 0.52 \end{array} \right\rangle & \left\langle \begin{array}{c} 0.588 \\ 0.412 \end{array} \right\rangle & \longrightarrow & \left\langle \begin{array}{c} 0.75 \\ 0.25 \end{array} \right\rangle \\ P(X_1) & P(X_2) & P(X_3) & P(X_4) & & P(X_\infty) \end{array}$$


- From yet another initial distribution $P(X_1)$:

$$\begin{array}{ccc} \left\langle \begin{array}{c} p \\ 1-p \end{array} \right\rangle & \dots & \longrightarrow \left\langle \begin{array}{c} 0.75 \\ 0.25 \end{array} \right\rangle \\ P(X_1) & & P(X_\infty) \end{array}$$

Video of Demo Ghostbusters Basic Dynamics



Video of Demo Ghostbusters Circular Dynamics



Stationary Distributions

- For most chains:

- Influence of the initial distribution gets less and less over time.
- The distribution we end up in is independent of the initial distribution

- Stationary distribution:


- The distribution we end up with is called the **stationary distribution** P_∞ of the chain
- It satisfies

$$P_\infty(X) = P_{\infty+1}(X) = \sum_x P(X|x)P_\infty(x)$$



Example: Stationary Distributions

- Question: What's $P(X)$ at time $t = \text{infinity}$?



$$P_\infty(\text{sun}) = P(\text{sun}|\text{sun})P_\infty(\text{sun}) + P(\text{sun}|\text{rain})P_\infty(\text{rain})$$


$$P_\infty(\text{rain}) = P(\text{rain}|\text{sun})P_\infty(\text{sun}) + P(\text{rain}|\text{rain})P_\infty(\text{rain})$$

$$P_\infty(\text{sun}) = 0.9P_\infty(\text{sun}) + 0.3P_\infty(\text{rain})$$

$$P_\infty(\text{rain}) = 0.1P_\infty(\text{sun}) + 0.7P_\infty(\text{rain})$$


$$P_\infty(\text{sun}) = 3P_\infty(\text{rain})$$

$$\text{Also: } P_\infty(\text{sun}) + P_\infty(\text{rain}) = 1$$



$$P_\infty(\text{sun}) = 3/4$$

$$P_\infty(\text{rain}) = 1/4$$




| X_{t-1} | X_t | $P(X_t X_{t-1})$ |
|-----------|-------|------------------|
| sun | sun | 0.9 |
| sun | rain | 0.1 |
| rain | sun | 0.3 |
| rain | rain | 0.7 |

- Alternatively: run simulation for a long (ideally infinite) time

Example: Stationary Distributions

- Question: What's $P(X)$ at time $t = \text{infinity}$?




- Matrix version

$$P_\infty = \begin{bmatrix} 0.9 & 0.3 \\ 0.1 & 0.7 \end{bmatrix} P_\infty$$

Shows that P_∞ is an eigenvector.

Also:


$$P_T = \begin{bmatrix} 0.9 & 0.3 \\ 0.1 & 0.7 \end{bmatrix}^T P_0$$



| X_{t-1} | X_t | $P(X_t X_{t-1})$ |
|-----------|-------|--------------------|
| sun | sun | 0.9 |
| sun | rain | 0.1 |
| rain | sun | 0.3 |
| rain | rain | 0.7 |

Example: Stationary Distributions

- Question: What's $P(X)$ at time $t = \text{infinity}$?




- Matrix version

$$P_\infty = \begin{bmatrix} 0.9 & 0.3 \\ 0.1 & 0.7 \end{bmatrix} P_\infty$$

Shows that P_∞ is an eigenvector. Also:

$$P_\infty = \begin{bmatrix} 0.9 & 0.3 \\ 0.1 & 0.7 \end{bmatrix}^\infty P_0 = \begin{bmatrix} 0.75 & 0.75 \\ 0.25 & 0.25 \end{bmatrix} P_0$$




| X_{t-1} | X_t | $P(X_t X_{t-1})$ |
|-----------|-------|--------------------|
| sun | sun | 0.9 |
| sun | rain | 0.1 |
| rain | sun | 0.3 |
| rain | rain | 0.7 |

Application of Stationary Distribution: Web Link Analysis


■ PageRank over a web graph

- Each web page is a state
- Initial distribution: uniform over pages
- Transitions:
 - With prob. c , uniform jump to a random page (dotted lines, not all shown)
 - With prob. $1-c$, follow a random outlink (solid lines)



■ Stationary distribution

- Will spend more time on highly reachable pages
- E.g. many ways to get to the Acrobat Reader download page
- Somewhat robust to link spam
- Google 1.0 returned the set of pages containing all your keywords in decreasing rank, now all search engines use link analysis along with many other factors (rank actually getting less important over time)




Hidden Markov Models




Hidden Markov Models

- Markov chains OK for games, weak for real robots




Hidden Markov Models


- Markov chains OK for games, weak for real robots



- Need observations to update your beliefs
- Hidden Markov models (HMMs)
 - Underlying Markov chain over states X
 - You observe **outputs (effects)** at each time step



Example: Weather HMM



- An HMM is defined by:

- Initial distribution: $P(X_1)$
 - Transitions: $P(X_t | X_{t-1})$
 - Emissions: $P(E_t | X_t)$

Transitions


| R_{t-1} | R_t | $P(R_t R_{t-1})$ |
|-----------|-------|--------------------|
| +r | +r | 0.7 |
| +r | -r | 0.3 |
| -r | +r | 0.3 |
| -r | -r | 0.7 |

Emissions

| R_t | U_t | $P(U_t R_t)$ |
|-------|-------|----------------|
| +r | +u | 0.9 |
| +r | -u | 0.1 |
| -r | +u | 0.2 |
| -r | -u | 0.8 |


Example: Ghostbusters HMM

- $P(X_1)$ = uniform
- $P(X' | X)$ = usually move clockwise, but sometimes move in a random direction or stay in place
- $P(R_{ij} | X)$ = same sensor model as before: red means close, green means far away.




| | | |
|-----|-----|-----|
| 1/9 | 1/9 | 1/9 |
| 1/9 | 1/9 | 1/9 |
| 1/9 | 1/9 | 1/9 |

$P(X_1)$




| | | |
|-----|-----|-----|
| 1/6 | 1/6 | 1/2 |
| 0 | 1/6 | 0 |
| 0 | 0 | 0 |

$P(X' | X = \langle 1, 2 \rangle)$




Video of Demo Ghostbusters – Circular Dynamics -- HMM



Conditional Independence


- HMMs have two important independence properties:
 - Markov hidden process: future depends on past via the present
 - Current observation independent of all else given current state



- Does this mean that evidence variables are guaranteed to be independent?

Conditional Independence


- HMMs have two important independence properties:
 - Markov hidden process: future depends on past via the present
 - Current observation independent of all else given current state




- Does this mean that evidence variables are guaranteed to be independent?
 - No, they are correlated by the hidden state

Real HMM Examples

- **Speech recognition HMMs:**
 - Observations are acoustic signals (continuous valued)
 - States are specific positions in specific words (so, tens of thousands)
- **Machine translation HMMs:**
 - Observations are words (tens of thousands)
 - States are translation options
- **Robot tracking:**
 - Observations are range readings (continuous)
 - States are positions on a map (continuous)




Inference in HMMs: Filtering



- Filtering, or monitoring, is the task of tracking the distribution $B_t(X) = P_t(X_t | e_1, \dots, e_t)$ (the belief state) over time
- We start with $B_1(X)$ in an initial setting, usually uniform
- As time passes, or we get observations, we update $B(X)$
- The Kalman filter was invented in the 60's and first implemented as a method of trajectory estimation for the Apollo program

Example: Robot Localization

Example from
Michael Pfeiffer




Prob

0

1



t=0



Sensor model: can read in which directions there is a wall,
never more than 1 mistake


Motion model: may not execute action with small prob.

Example: Robot Localization



Lighter grey: was possible to get the reading, but less likely b/c required 1 mistake

Example: Robot Localization




Prob


0

1

t=2



Example: Robot Localization




Prob


0

1

t=3



Example: Robot Localization




Prob


0

1

$t=4$



Example: Robot Localization




Prob

0

1

t=5




HMM Inference: Find State Given Evidence

- We are given evidence at each time and want to know


$$B_t(X) = P(X_t | e_{1:t})$$

- Idea: start with $P(X_1)$ and derive $B_t(X)$ in terms of $B_{t-1}(X)$
 - Two steps: **Passage of Time** & **Observation**




Inference: Base Cases


Passage of Time:



Observation:



Passage of Time: Base Case



Have: $P(X_1)$ $P(X_2|X_1)$


Want: $P(X_2)$

$$\begin{aligned} P(x_2) &= \sum_{x_1} P(x_1, x_2) \\ &= \sum_{x_1} P(x_1)P(x_2|x_1) \end{aligned}$$

Passage of Time: General Case

- Assume we have current belief $P(X \mid \text{evidence to date})$

$$B(X_t) = P(X_t | e_{1:t})$$



- Then, after one time step passes:

$$\begin{aligned} P(X_{t+1} | e_{1:t}) &= \sum_{x_t} P(X_{t+1}, x_t | e_{1:t}) \\ &= \sum_{x_t} P(X_{t+1} | x_t, e_{1:t}) P(x_t | e_{1:t}) \\ &= \sum_{x_t} P(X_{t+1} | x_t) P(x_t | e_{1:t}) \end{aligned}$$

- Or compactly:

$$B'(X_{t+1}) = \sum_{x_t} P(X' | x_t) B(x_t)$$

- Basic idea: beliefs get “pushed” through the transitions
 - With the “B” notation, we have to be careful about what time step t the belief is about, and what evidence it includes

Example: Passage of Time

- As time passes, uncertainty “accumulates”

(Transition model: ghosts usually go clockwise)

| | | | | | |
|-------|-------|-------|-------|-------|-------|
| <0.01 | <0.01 | <0.01 | <0.01 | <0.01 | <0.01 |
| <0.01 | <0.01 | <0.01 | <0.01 | <0.01 | <0.01 |
| <0.01 | <0.01 | 1.00 | <0.01 | <0.01 | <0.01 |
| <0.01 | <0.01 | <0.01 | <0.01 | <0.01 | <0.01 |



T = 1

| | | | | | |
|-------|-------|-------|-------|-------|-------|
| <0.01 | <0.01 | <0.01 | <0.01 | <0.01 | <0.01 |
| <0.01 | <0.01 | 0.06 | <0.01 | <0.01 | <0.01 |
| <0.01 | 0.76 | 0.06 | 0.06 | <0.01 | <0.01 |
| <0.01 | <0.01 | 0.06 | <0.01 | <0.01 | <0.01 |


T = 2

| | | | | | |
|------|------|-------|-------|-------|-------|
| 0.05 | 0.01 | 0.05 | <0.01 | <0.01 | <0.01 |
| 0.02 | 0.14 | 0.11 | 0.35 | <0.01 | <0.01 |
| 0.07 | 0.03 | 0.05 | <0.01 | 0.03 | <0.01 |
| 0.03 | 0.03 | <0.01 | <0.01 | <0.01 | <0.01 |

T = 5



Observation: Base Case



Have: $P(X_1)$ $P(E_1|X_1)$

Want: $P(X_1|e_1)$

$$\begin{aligned} P(x_1|e_1) &= P(x_1, e_1)/P(e_1) \\ &\propto_{X_1} P(x_1, e_1) \\ &= P(x_1)P(e_1|x_1) \end{aligned}$$

Also can write as:

$$P(x_1|e_1) = \frac{P(x_1)P(e_1|x_1)}{\sum_{x'} P(x')P(e_1|x')}$$

Observation: General Case

- Assume we have current belief $P(X \mid \text{previous evidence})$:


$$B'(X_{t+1}) = P(X_{t+1} | e_{1:t})$$

- Then, after evidence comes in:

$$\begin{aligned} P(X_{t+1} | e_{1:t+1}) &= P(X_{t+1}, e_{t+1} | e_{1:t}) / P(e_{t+1} | e_{1:t}) \\ &\propto_{X_{t+1}} P(X_{t+1}, e_{t+1} | e_{1:t}) \\ &= P(e_{t+1} | e_{1:t}, X_{t+1}) P(X_{t+1} | e_{1:t}) \\ &= P(e_{t+1} | X_{t+1}) P(X_{t+1} | e_{1:t}) \end{aligned}$$

- Or, compactly:

$$B(X_{t+1}) \propto_{X_{t+1}} P(e_{t+1} | X_{t+1}) B'(X_{t+1})$$



- Basic idea: beliefs “reweighted” by likelihood of evidence
- Unlike passage of time, we have to renormalize

Example: Observation

- As we get observations, beliefs get reweighted, uncertainty “decreases”


| | | | | | |
|------|------|-------|-------|-------|-------|
| 0.05 | 0.01 | 0.05 | <0.01 | <0.01 | <0.01 |
| 0.02 | 0.14 | 0.11 | 0.35 | <0.01 | <0.01 |
| 0.07 | 0.03 | 0.05 | <0.01 | 0.03 | <0.01 |
| 0.03 | 0.03 | <0.01 | <0.01 | <0.01 | <0.01 |

Before observation


| | | | | | |
|-------|-------|-------|-------|-------|-------|
| <0.01 | <0.01 | <0.01 | <0.01 | 0.02 | <0.01 |
| <0.01 | <0.01 | <0.01 | 0.83 | 0.02 | <0.01 |
| <0.01 | <0.01 | 0.11 | <0.01 | <0.01 | <0.01 |
| <0.01 | <0.01 | <0.01 | <0.01 | <0.01 | <0.01 |

After observation

$$B(X) \propto P(e|X)B'(X)$$




Two Steps: Passage of Time + Observation



Pacman – Sonar



Video of Demo Pacman – Sonar (with beliefs)



Example: Weather HMM



Passage of Time:


$$B'(X_{t+1}) = \sum_{x_t} P(X_{t+1}|x_t) B(x_t)$$

Observation:

$$B(X_{t+1}) \propto_{X_{t+1}} P(e_{t+1}|X_{t+1}) B'(X_{t+1})$$

$B(+r) = 0.5$
 $B(-r) = 0.5$

$B'(+r) = ?$
 $B'(-r) = ?$



$P(X_{t+1}|X_t)$

| R_t | R_{t+1} | $P(R_{t+1} R_t)$ |
|-------|-----------|------------------|
| +r | +r | 0.7 |
| +r | -r | 0.3 |
| -r | +r | 0.3 |
| -r | -r | 0.7 |

$P(E_t|X_t)$

| R_t | U_t | $P(U_t R_t)$ |
|-------|-------|--------------|
| +r | +u | 0.9 |
| +r | -u | 0.1 |
| -r | +u | 0.2 |
| -r | -u | 0.8 |

Example: Weather HMM



Passage of Time:

$$B'(X_{t+1}) = \sum_{x_t} P(X_{t+1}|x_t) B(x_t)$$

Observation:


$$B(X_{t+1}) \propto_{X_{t+1}} P(e_{t+1}|X_{t+1}) B'(X_{t+1})$$

$$B'(+r) = 0.5 \cdot 0.7 + 0.5 \cdot 0.3 = 0.5$$

$$B'(-r) = 0.5 \cdot 0.3 + 0.5 \cdot 0.7 = 0.5$$

$$B(+r) = 0.5$$

$$B(-r) = 0.5$$




$P(X_{t+1}|X_t)$

| R_t | R_{t+1} | $P(R_{t+1} R_t)$ |
|-------|-----------|------------------|
| +r | +r | 0.7 |
| +r | -r | 0.3 |
| -r | +r | 0.3 |
| -r | -r | 0.7 |

$P(E_t|X_t)$

| R_t | U_t | $P(U_t R_t)$ |
|-------|-------|--------------|
| +r | +u | 0.9 |
| +r | -u | 0.1 |
| -r | +u | 0.2 |
| -r | -u | 0.8 |

Example: Weather HMM




Passage of Time:

$$B'(X_{t+1}) = \sum_{x_t} P(X_{t+1}|x_t) B(x_t)$$

Observation:

$$B(X_{t+1}) \propto_{X_{t+1}} P(e_{t+1}|X_{t+1}) B'(X_{t+1})$$

$$\begin{array}{l} B(+r) = 0.5 \\ B(-r) = 0.5 \end{array} \quad \begin{array}{l} \nearrow \\ \\ \searrow \end{array} \quad \begin{array}{l} B'(+r) = 0.5 \\ B'(-r) = 0.5 \\ \downarrow \\ B(+r) = ? \\ B(-r) = ? \end{array}$$





$P(X_{t+1}|X_t)$

| R_t | R_{t+1} | $P(R_{t+1} R_t)$ |
|-------|-----------|------------------|
| +r | +r | 0.7 |
| +r | -r | 0.3 |
| -r | +r | 0.3 |
| -r | -r | 0.7 |

$P(E_t|X_t)$

| R_t | U_t | $P(U_t R_t)$ |
|-------|-------|--------------|
| +r | +u | 0.9 |
| +r | -u | 0.1 |
| -r | +u | 0.2 |
| -r | -u | 0.8 |

Example: Weather HMM




Passage of Time:

$$B'(X_{t+1}) = \sum_{x_t} P(X_{t+1}|x_t) B(x_t)$$

Observation:

$$B(X_{t+1}) \propto_{X_{t+1}} P(e_{t+1}|X_{t+1}) B'(X_{t+1})$$





$P(X_{t+1}|X_t)$

| R_t | R_{t+1} | $P(R_{t+1} R_t)$ |
|-------|-----------|------------------|
| +r | +r | 0.7 |
| +r | -r | 0.3 |
| -r | +r | 0.3 |
| -r | -r | 0.7 |

$P(E_t|X_t)$

| R_t | U_t | $P(U_t R_t)$ |
|-------|-------|--------------|
| +r | +u | 0.9 |
| +r | -u | 0.1 |
| -r | +u | 0.2 |
| -r | -u | 0.8 |

Example: Weather HMM




Passage of Time:

$$B'(X_{t+1}) = \sum_{x_t} P(X_{t+1}|x_t) B(x_t)$$

Observation:

$$B(X_{t+1}) \propto_{X_{t+1}} P(e_{t+1}|X_{t+1}) B'(X_{t+1})$$



$P(X_{t+1}|X_t)$

| R_t | R_{t+1} | $P(R_{t+1} R_t)$ |
|-------|-----------|------------------|
| +r | +r | 0.7 |
| +r | -r | 0.3 |
| -r | +r | 0.3 |
| -r | -r | 0.7 |

$P(E_t|X_t)$

| R_t | U_t | $P(U_t R_t)$ |
|-------|-------|--------------|
| +r | +u | 0.9 |
| +r | -u | 0.1 |
| -r | +u | 0.2 |
| -r | -u | 0.8 |

What we did today

- **Markov Chains & their Stationary Distributions**
 - How beliefs about state change with passage of time
- **Hidden Markov Models (HMMs) formulation**
 - How beliefs change with passage of time and evidence
- **Filtering with HMMs**
 - How to infer beliefs from evidence

Next Time: More Filtering!
