


Q1. Bayes Nets: Variable Elimination



	$P(A)$
$+a$	0.25
$-a$	0.75

	$P(B A)$	$+b$	$-b$
$+a$		0.5	0.5
$-a$		0.25	0.75

	$P(C A)$	$+c$	$-c$
$+a$		0.2	0.8
$-a$		0.6	0.4


	$P(D B)$	$+d$	$-d$
$+b$		0.6	0.4
$-b$		0.8	0.2

	$P(E B)$	$+e$	$-e$
$+b$		0.25	0.75
$-b$		0.1	0.9

- (a) Using the Bayes' Net and conditional probability tables above, calculate the following quantities:
- (i) $P(+b | +a) =$
 - (ii) $P(+a, +b) =$
 - (iii) $P(+a | +b) =$
- (b) Now we are going to consider variable elimination in the Bayes' Net above.
- (i) Assume we have the evidence $+c$ and wish to calculate $P(E | +c)$. What factors do we have initially?
 - (ii) If we eliminate variable B, we create a new factor. What probability does that factor correspond to?
 - (iii) What is the equation to calculate the factor we create when eliminating variable B?
 - (iv) After eliminating variable B, what are the new set of factors? As in (ii), write the probabilities that the factors represent. For each factor, also provide its size.
 - (v) Now assume we have the evidence $-c$ and are trying to calculate $P(A | -c)$. What is the most efficient elimination ordering? If more than one ordering is most efficient, provide any one of them.
 - (vi) Once we have run variable elimination and have $f(A, -c)$ how do we calculate $P(+a | -c)$?


Q2. Bayes Nets and Joint Distributions

- (a) Write down the joint probability distribution associated with the following Bayes Net. Express the answer as a product of terms representing individual conditional probabilities tables associated with this Bayes Net:



- (b) Draw the Bayes net associated with the following joint distribution:

$$P(A) \cdot P(B) \cdot P(C|A, B) \cdot P(D|C) \cdot P(E|B, C)$$



- (c) Do the following products of factors correspond to a valid joint distribution over the variables A, B, C, D ? (Circle FALSE or TRUE.)

(i) FALSE TRUE $P(A) \cdot P(B) \cdot P(C|A) \cdot P(C|B) \cdot P(D|C)$

(ii) FALSE TRUE $P(A) \cdot P(B|A) \cdot P(C) \cdot P(D|B, C)$

(iii) FALSE TRUE $P(A) \cdot P(B|A) \cdot P(C) \cdot P(C|A) \cdot P(D)$

(iv) FALSE TRUE $P(A|B) \cdot P(B|C) \cdot P(C|D) \cdot P(D|A)$


(d) What factor can be multiplied with the following factors to form a valid joint distribution? (Write “none” if the given set of factors can’t be turned into a joint by the inclusion of exactly one more factor.)

(i) $P(A) \cdot P(B|A) \cdot P(C|A) \cdot P(E|B, C, D)$

(ii) $P(D) \cdot P(B) \cdot P(C|D, B) \cdot P(E|C, D, A)$

(e) Answer the next questions based off of the Bayes Net below:

All variables have domains of $\{-1, 0, 1\}$



(i) Before eliminating any variables or including any evidence, how many entries does the factor at G have?

(ii) Now we observe $e = 1$ and want to query $P(D|e = 1)$, and you get to pick the first variable to be eliminated.

- Which choice would create the **largest** factor f_1 ?

- Which choice would create the **smallest** factor f_1 ?

Q3. Probability and Bayes Nets

- (a) A, B, and C are random variables with binary domains. How many entries are in the following probability tables and what is the sum of the values in each table? Write a “?” in the box if there is not enough information given.

Table	Size	Sum
$P(A, B C)$		
$P(A +b,+c)$		
$P(+a B)$		

- (b) Circle true if the following probability equalities are valid and circle false if they are invalid (leave it blank if you don't wish to risk a guess). Each True/False question is worth 1 points. Leaving a question blank is worth 0 points. **Answering incorrectly is worth -1 points.**

No independence assumptions are made.

(i) [true or false] $P(A, B) = P(A|B)P(A)$

(ii) [true or false] $P(A|B)P(C|B) = P(A, C|B)$

(iii) [true or false] $P(B, C) = \sum_{a \in A} P(B, C|A)$

(iv) [true or false] $P(A, B, C, D) = P(C)P(D|C)P(A|C, D)P(B|A, C, D)$

- (c) Space Complexity of Bayes Nets

Consider a joint distribution over N variables. Let k be the domain size for all of these variables, and let d be the maximum indegree of any node in a Bayes net that encodes this distribution.

(i) What is the space complexity of storing the entire joint distribution? Give an answer of the form $O(\cdot)$.

(ii) Draw an example of a Bayes net over four binary variables such that it takes less space to store the Bayes net than to store the joint distribution.

(iii) Draw an example of a Bayes net over four binary variables such that it takes more space to store the Bayes net than to store the joint distribution.