
Final

Last Name	First Name	SID
-----------	------------	-----

Rules.

- **Unless otherwise stated, all your answers need to be justified and your work must be shown. Answers without sufficient justification will get no credit.**
- You have 160 minutes to complete the exam and 10 minutes exclusively for submitting your exam to Gradescope. (DSP students with $X\%$ time accomodation should spend $160 \cdot X\%$ time on the exam and 10 minutes to submit).
- Collaboration with others is strictly prohibited.
- You may reference your notes, the textbook, and any material that can be found through the course website. You may use Google to search up general knowledge. However, searching up a question is not allowed.
- You may not use online solvers or graphing tools (ex. WolframAlpha, Desmos, Python). Simple functions (ex. combinations, multiplication) are fine.
- For any clarifications you have, please create a private Piazza post. We will have a Google Doc that shows our official clarifications.

Problem	points earned	out of
Problem 1		16
Problem 2		7
Problem 3		7
Problem 4		7
Problem 5		8
Problem 6		10
Problem 7		10
Problem 8		13
Problem 9		7
Problem 10		8
Problem 11		7
Total		100

1 True or False (4 + 4 + 4 + 4 points)

For each of the following, say whether the assertion is true or false. If it is true, provide justification, and if it is false, give a counterexample.

(a) For any finite sample space Ω and any event $A \subseteq \Omega$, $\Pr(A) = \frac{|A|}{|\Omega|}$.

(b) If $Y = X + Z$ is Gaussian, then X and Z are both marginally Gaussian.

(c) Michael and Kevin are playing a game where Michael scores $X_i \sim \text{Geometric}(1/m)$ points and Kevin scores $Y_i \sim \text{Geometric}(1/k)$ points at round i , independently of other rounds. Then, as the number of rounds goes to infinity, the average total number of points they score per round converges almost surely to $m + k$.

(d) Suppose a random variable X is bounded in $[0, 1]$, and furthermore suppose that $\mathbb{E}[X] \geq \epsilon$. Then $\Pr(X \geq \epsilon/2) \geq \epsilon/2$.

2 Waiting Game (7 points)

The amount of time you have to wait for UC Berkeley to announce the PNP policy is exponentially distributed with parameter λ . Solve for the optimal Chernoff upper bound (by applying Markov's inequality to e^{sX}) for the probability that you have to wait for longer than some constant a .

3 Terms and Conditionings (7 points)

Suppose $X \sim \min\{q, Y\}$, where $Y \sim \text{Geom}(p)$ is a geometric RV with parameter p and q is a positive integer. Calculate $\mathbb{E}[X]$ in terms of p and q .

4 Homework Party (7 points)

Michael is trying to budget time for his problem set. He knows that the number of problems on the problem set is Poisson(6) distributed, and the probability that he can solve any given problem is $\frac{1}{4}$, and is independent of all other problems. If he can solve the problem, the amount of time he spends on the problem is Exponential ($\frac{1}{6}$) distributed; otherwise, the amount of time he spends on the problem is Exponential ($\frac{1}{10}$) distributed. What is the expected amount of time that Michael will spend on the problem set?

5 Seasonal Depression (8 points)

Kevin is a store owner in a part of town with strange weather. The weather alternates between two states: sunny and rainy. The length of each sunny period and rainy period is determined by an independent exponential with rate 1.

- If the weather is sunny, customers arrive at a store according to a Poisson process with rate 1 and each one leaves with rate 1.
- If the weather is rainy, customers arrive at a store according to a Poisson process with rate 1 and do not leave.

Suppose it is currently sunny and there are no customers in the store. What is the expected amount of time before there are at least 2 people in the store?

6 Geometric Perspective on Variance (3 + 5 + 2 points)

In this problem, we prove the identity $\text{var}(X) = \mathbb{E}[\text{var}(X|Y)] + \text{var}(\mathbb{E}[X|Y])$. Assume X and Y are zero mean. You may use that $\text{var}(X|Y) = \mathbb{E}[(X - \mathbb{E}[X|Y])^2|Y]$.

(a) First, show that $\mathbb{E}[\text{var}(X|Y)] = \text{var}(X - \mathbb{E}[X|Y])$.

(b) Draw X , $\mathbb{E}[X|Y]$, and $X - \mathbb{E}[X|Y]$ in the Hilbert space of random variables. Specify and justify the angle between $\mathbb{E}[X|Y]$ and $X - \mathbb{E}[X|Y]$.

(c) Use the previous two parts to conclude the identity $\text{var}(X) = \mathbb{E}[\text{var}(X|Y)] + \text{var}(\mathbb{E}[X|Y])$.

8 Hidden Markovs Among Us (3 + 3 + 3 + 2 + 2 points)

Let the number of people infected by COVID on day n be denoted by X_n . Each day, X_n increases by 1 with probability $\frac{2}{3}$ or decreases by 1 with probability $\frac{1}{3}$. If $X_n = 0$, it stays the same with probability $\frac{1}{3}$ or increases with probability $\frac{2}{3}$. Then, let $Y_n \sim \text{Binomial}(X_n, \frac{3}{4})$ represent the number of people who test positive for COVID, i.e. that we report having COVID. Assume $X_0 = 1$.

(a) What is the MAP estimate of X_2 given that we observe $Y_2 = 1$?

(b) What is the MLE estimate of X_2 given $Y_2 = 1$? Multiple values are fine.

(c) What is the LLSE estimator of X_2 given $Y_2 = 1$? (You may use $\text{Var}(Y_2) \approx 1.056$)

(d) What is the MMSE estimator of X_2 given $Y_2 = 1$?

(e) Is the Markov chain $\{X_t\}_{t=0}^{\infty}$ positive recurrent, null recurrent, or transient?

9 Dungeons and Dragons (7 points)

In a game of Dungeons and Dragons, Aditya suspects Catherine is using a loaded 20-sided die. However, he doesn't want to risk falsely accusing her, so he conducts a hypothesis test to upper-bound the probability of a false alarm. Suppose if $X = 0$, the die is fair and a roll Y is distributed according to $\Pr(Y = y|X = 0) = 0.05$ for $1 \leq y \leq 20$. If $X = 1$, the die is loaded, and rolls have the distribution

$$\Pr(Y = y|X = 1) = \begin{cases} 0.025 & 1 \leq y \leq 10 \\ 0.075 & 11 \leq y \leq 20 \end{cases}$$

(or more simply, the probability of being greater than 10 is three times the probability of being less than or equal to 10). Construct a Neyman-Pearson decision rule to maximize the probability Aditya is correct if he accuses Catherine of cheating, while constraining the probability Aditya falsely accuses Catherine to be ≤ 0.05 .

10 Delayed Kalman Filter (6 + 2 points)

Consider a process with the transition rule $x_{n+1} = ax_n + v_n$ where $v_n \sim \mathcal{N}(0, \sigma_v^2)$. We can only observe the process at even-numbered times, i.e. we see $y_{2n} = x_{2n} + w_{2n}$, where $w_n \sim \mathcal{N}(0, \sigma_w^2)$.

1. Find a recurrence relation for the MMSE of the even states $\hat{x}_{2n} = \mathbb{E}[x_{2n}|y_0, y_2, \dots, y_{2n}]$ in terms of \hat{x}_{2n-2} .

2. Find a recurrence relation for the MMSE of the odd states $\hat{x}_{2n+1} = \mathbb{E}[x_{2n+1}|y_0, y_2, \dots, y_{2n}]$ in terms of \hat{x}_{2n} .

