
Midterm 1

Last Name	First Name	SID
-----------	------------	-----

Rules.

- **Unless otherwise stated, all your answers need to be justified and your work must be shown. Answers without sufficient justification will get no credit.**
- You have 70 minutes to complete the exam and 10 minutes exclusively for submitting your exam to Gradescope. (DSP students with $X\%$ time accomodation should spend $70 \cdot X\%$ time on the exam and 10 minutes to submit).
- Collaboration with others is strictly prohibited.
- You may reference your notes, the textbook, and any material that can be found through the course website. You may use Google to search up general knowledge. However, searching up a question is not allowed.
- You may not use online solvers or graphing tools (ex. WolframAlpha, Desmos, Python). Simple functions (ex. combinations, multiplication) are fine.
- For any clarifications you have, please create a private Piazza post. We will have a Google Doc that shows our official clarifications.

Problem	points earned	out of
Honor Code		1
Problem 1		60
Problem 2		27
Problem 3		14
Total		102

1 Assorted Problems

(a) **Flickering Lights [10 points]**

Suppose there are n lights that turn on at the same time, each of which have a lifespan modeled by an exponential distribution with parameter λ . Let t_1, t_2, \dots, t_{n-1} describe the amount of time elapsed between successive failures, where t_i is the time between the i -th failure and the $i + 1$ -th failure. Find the expected value of the minimum of all the t_i ($1 \leq i \leq n - 1$).

Example: Say we have 4 lights that shutoff at times 2, 4, 7, and 8. The times elapsed between shutoffs are 2, 3, and 1.

(b) **Unfortunate Occurrences [10 points]**

Suppose that on any given day there is a wildfire with probability p , and all the days are independent. Given that there is at least one wildfire within the next n days, what is the expected value of the number of days until the next wildfire (i.e. a value between 1 and n possibly dependent on p and/or n)?

(c) **Be Reasonable [10 points]**

Let's say that for some real $b \geq 1$ that a random variable X is b -reasonable if it satisfies:

$$\mathbb{E}[X^4] \leq b(\mathbb{E}[X^2])^2$$

Suppose X is b -reasonable and that $\mathbb{E}[X] = 0$ and $\text{var}(X) = \sigma^2$. Prove that

$$\Pr[|X| \geq t\sigma] \leq \frac{b}{t^4}$$

(d) **Normal Product Sequence [10 points]**

Let X_1 have some distribution whose moments (i.e. $\mathbb{E}[X_1], \mathbb{E}[X_1^2], \mathbb{E}[X_1^3], \dots$) are all known. For $i \geq 2$, $X_i \sim \mathcal{N}(X_1 X_2 \dots X_{i-1}, 1)$, i.e. a normal distribution with mean equal to the product of the previous elements of the sequence. Find the expectation of X_4 in terms of moments of X_1 .

(e) **Pascal's Coins [10 points]**

Someone tells you that they flipped $n \geq 1$ fair coins and got 4 heads, but they don't tell you what n is. What is the smallest integer MLE estimate of n ?

(f) **Predetermined Fate [10 points]**

Let Z_i equal i^4 with probability $\frac{1}{i^2}$ and -1 with probability $1 - \frac{1}{i^2}$, and let $W_n = \sum_{i=1}^n Z_i$. Show that the probability of $W_n \rightarrow -\infty$ as $n \rightarrow \infty$ is one. (Hint: $\sum_{i=1}^{\infty} \frac{1}{i^2} = \frac{\pi^2}{6}$)

2 Chess Antics

(a) Knight's Journey [10 points]

Consider a 13×13 chess board. An “upward knight”, starting in the bottom left corner $(1, 1)$, tries to move to the upper right corner $(13, 13)$. This knight can only make one of two moves:

- (a) It can move up by two, right by one: $(x, y) \rightarrow (x + 1, y + 2)$
- (b) It can move up by one, right by two: $(x, y) \rightarrow (x + 2, y + 1)$

If both moves are available (would not take the knight out of bounds), it chooses one uniformly at random, or else it takes the only available move. What is the probability the knight travels through the center square $(7, 7)$? (**Hint:** think how many of each type of move is required to reach $(13, 13)$ or $(7, 7)$)

(b) Queen's Journey [10 points]

Suppose we have a queen on a 3×3 chess board. The queen's initial position is random, and it proceeds to make random legal moves (the queen can move horizontally, vertically, or diagonally for any number of squares). Assume the queen never stays in the same square. What is the fraction of time the queen spends in any of the four corners of the board as its number of moves approaches infinity? (**Hint:** can this be modeled as a Markov chain?)


(c) Final Showdown [7 points]

Kevin and Michael are playing a chess match consisting of many games. Let K equal the number of points Kevin has won minus the number of points Michael has won. Both players are equally matched, so Kevin and Michael are both equally likely to win each game. If Kevin wins a game, Kevin gets a point; if Michael wins, Michael gets a point; there are no draws. The match ends when $K = 5$ (Kevin wins) or when $K = -5$ (Michael wins).

- (a) Let K_t denote the value of K after t games have been played and let $K_{t+1} = K_t$ if the match has already ended by time t . Draw K_t as a Markov chain and find the corresponding transition matrix P (such that $\pi_{t+1} = P\pi_t$ models the distribution of K_{t+1}).
- (b) Is the Markov chain irreducible? What are the states that have can nonzero probability in the stationary/invariant distribution?

3 Graphical Density

Two random variables X and Y have a joint density as pictured below:



Note that the formula for a circle of radius 1 centered at the origin is $x^2 + y^2 = 1$, and the right portion of the graph is just one quarter of a unit circle.

- What is A ? [2 points]
- Find the marginals $f_X(x)$ and $f_Y(y)$. Write your answer in terms of A . [4 points]
- What is $f_{Y|X}(y|x)$? [2 points]
- What is $\mathbb{E}[Y|X]$? [2 points]
- What is $\mathbb{E}[Y]$? Write your final answer in terms of A . [4 points]