

Discussion 2

Spring 2020

1. Sampling without Replacement

Suppose you have N items, G of which are good and B of which are bad (B , G , and N are positive integers, $B + G = N$). You start to draw items without replacement, and suppose that the first good item appears on draw X . Compute the mean and variance of X .

2. Poisson Merging

Let X and Y be independent Poisson random variables with means λ and μ respectively. Prove that $X + Y \sim \text{Poisson}(\lambda + \mu)$. (This is known as **Poisson merging**.) Note that it is **not** sufficient to use linearity of expectation to say that $X + Y$ has mean $\lambda + \mu$. You are asked to prove that the *distribution* of $X + Y$ is Poisson.

Note: You may need to use the Binomial theorem: $(x + y)^n = \sum_{i=0}^n \binom{n}{i} x^i y^{n-i}$. This poisson merging property will be extensively used when we discuss Poisson processes.

3. Clustering Coefficient

This problem will explore an important probabilistic concept of clustering that is widely used in machine learning applications today. Consider n students, where n is a positive integer. For each pair of students $i, j \in \{1, \dots, n\}$, $i \neq j$, they are friends with probability p , independently of other pairs. We assume that friendship is mutual. We can see that the friendship among the n students can be represented by an undirected graph G . Let $N(i)$ be the number of friends of student i and $T(i)$ be the number of triangles attached to student i . We define the **clustering coefficient** $C(i)$ for student i as follows:

$$C(i) = \frac{T(i)}{\binom{N(i)}{2}}.$$




Figure 1: Friendship and clustering coefficient.

The clustering coefficient is not defined for the students who have no friends. An example is shown in Figure 1. Student 3 has 4 friends (1, 2, 4, 5) and there are two triangles attached to student 3, i.e., triangle 1-2-3 and triangle 2-3-4. Therefore $C(3) = 2/\binom{4}{2} = 1/3$.

Find $\mathbb{E}[C(i) \mid N(i) \geq 2]$.