
EECS 16A Designing Information Devices and Systems I Discussion 4A
 Fall 2019

1. Proofs

- (a) Prove the following statement (proved earlier in lecture): If the columns of A are linearly dependent, then $A\vec{x} = \vec{b}$ does not have a unique solution.
- (b) Often, when one is asked to prove something you are asked to prove something of the following nature:
- $P \implies Q$. This is read as P implies Q .

Identify P and Q in the theorem you just proved above.

There are a couple of things to remember when reading these statements. First, is that the direction of implication matters.

- If you prove $P \implies Q$, this does not mean that $Q \implies P$ is also true.

Suppose someone tells you that $P \implies Q$ is true. Then you find out later that Q is actually false. What can you say about P ?

- If $P \implies Q$ and Q is false, then P must be false.

2. Identifying a Basis

Does each of these sets of vectors describe a basis for \mathbb{R}^3 ? If the vectors do not form a basis for \mathbb{R}^3 , can they be thought of as a basis for some other vector space?

$$V_1 = \left\{ \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}, \begin{bmatrix} 1 \\ 0 \\ 1 \end{bmatrix} \right\} \quad V_2 = \left\{ \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix}, \begin{bmatrix} 1 \\ 0 \\ 1 \end{bmatrix}, \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix} \right\} \quad V_3 = \left\{ \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix}, \begin{bmatrix} 1 \\ 0 \\ 1 \end{bmatrix}, \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix} \right\}$$

3. Exploring Column Spaces and Null Spaces

- The **column space** is the **span** of the column vectors of the matrix.
- The **null space** is the set of input vectors that output the zero vector.

For the following matrices, answer the following questions:

- What is the column space of \mathbf{A} ? What is its dimension?
- What is the null space of \mathbf{A} ? What is its dimension?
- Are the column spaces of the row reduced matrix \mathbf{A} and the original matrix \mathbf{A} the same?
- Do the columns of \mathbf{A} form a basis for \mathbb{R}^2 ? Why or why not?

(a) $\begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix}$

(b) $\begin{bmatrix} 0 & 1 \\ 0 & 1 \end{bmatrix}$

$$(c) \begin{bmatrix} 1 & 2 \\ -1 & 1 \end{bmatrix}$$

$$(d) \begin{bmatrix} -2 & 4 \\ 3 & -6 \end{bmatrix}$$

$$(e) \begin{bmatrix} 1 & -1 & -2 & -4 \\ 1 & 1 & 3 & -3 \end{bmatrix}$$