


CAL CAMPUS



SOUTH OF DWIGHT

(b) Can you set this up as a system of equations? Is it linear? If it's not linear, can you think of a way to make it linear? Now, how do you set this up in matrix form?

Hint: Set (0,0) to be Channing and Bowditch.

Hint 2: Distance = $\sqrt{(x_1 - x_2)^2 + (y_1 - y_2)^2}$


Hint 3: You don't need all 4 equations. You have two unknowns, x and y . You know from lecture that you need three circles to uniquely find a point. How can you use the third circle/equation to get two equations and two unknowns?

(c) Suppose Mr. Muffin is moving fast, and by the time you get to destination in part (a) he's already run off! You check the logs of the cell towers again, and see the following updated messages:

Sensor	Distance
N	2.2
W	Out of Range
E	1.1
S	Out of Range

Can you find Mr. Muffin?

CAL CAMPUS




SOUTH OF DWIGHT

- (d) Mr. Muffin is a very mischievous puppy, and while playing and running around he damaged his collar. The transmitter on his collar will still send a signal to the receiver towers, but the distance sensor has noise. You check the logs of the cell towers, and they have received the following messages:

Sensor	Distance
N	1.7 ± 0.5
W	2.1 ± 0.2
E	Out of Range
S	Out of Range

On the map provided, identify where Mr. Muffin is! Can you find exactly where he is?

CAL CAMPUS



SOUTH OF DWIGHT

2. A review of Inner Products

Find the inner product of the following three pairs of vectors.

(a)

$$\begin{bmatrix} 1 \\ 0 \\ 1 \end{bmatrix}, \begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix}$$

(b)

$$\begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix}, \begin{bmatrix} 1 \\ 0 \\ 1 \end{bmatrix}$$

(c)

$$\begin{bmatrix} 1 \\ 0 \\ 3 \end{bmatrix}, \begin{bmatrix} -3 \\ 2 \\ 1 \end{bmatrix}$$

3. From Inner Products To Projections

Given that $\langle \vec{x}, \vec{y} \rangle$ is a measure of similarity between two vectors, let's try to use this to find how much of one vector \vec{y} is in the direction of another vector \vec{x} .

(a) Let's start with $\langle \vec{x}, \vec{y} \rangle$. We want a quantity that is independent of the norm of \vec{x} , $\|\vec{x}\|$. Is $\langle \vec{x}, \vec{y} \rangle$ independent of the norm? Consider $\langle \vec{x}, \vec{y} \rangle$ for the examples below.

$$\vec{x} = \begin{bmatrix} 1 \\ 1 \end{bmatrix}, \vec{y} = \begin{bmatrix} 2 \\ 1 \end{bmatrix}, \vec{x} = \begin{bmatrix} 2 \\ 2 \end{bmatrix}, \vec{y} = \begin{bmatrix} 2 \\ 1 \end{bmatrix}$$

(b) Suppose we divide $\langle \vec{x}, \vec{y} \rangle$ by the norm of \vec{x} , $\|\vec{x}\|$, to get $\frac{\langle \vec{x}, \vec{y} \rangle}{\|\vec{x}\|}$. Is this new quantity independent of the norm of \vec{x} ? Test it on the examples above.

(c) We now have a scalar quantity that represents how much of \vec{y} is in the direction of \vec{x} . Let's try to find a vector that is how much of \vec{y} is in the \vec{x} direction. That is, we are looking for a vector \vec{z} that has a norm of $\frac{\langle \vec{x}, \vec{y} \rangle}{\|\vec{x}\|}$ and points in the same direction as \vec{x} .

(d) Given the projection between two vectors, defined as $\text{proj}_{\vec{x}} \vec{y} = \frac{\langle \vec{x}, \vec{y} \rangle}{\|\vec{x}\|^2} \vec{x}$, prove the Cauchy-Schwarz inequality, $|\langle \vec{x}, \vec{y} \rangle| \leq \|\vec{x}\| \|\vec{y}\|$.

(e) Consider the quantity $\frac{\langle \vec{x}, \vec{y} \rangle}{\|\vec{x}\| \|\vec{y}\|}$. What is the maximum this quantity could be? When does this occur? What is the minimum this quantity could be? When does this occur?

(f) We define the angle between two vectors as $\cos(\theta) = \frac{\langle \vec{x}, \vec{y} \rangle}{\|\vec{x}\| \|\vec{y}\|}$. When do two vectors have an angle of 90° between them? When do they have an angle of 0° ? When do they have an angle of 180° ?