

---

EECS 16A    Designing Information Devices and Systems I  
Fall 2022    Discussion 7A

---

**1. Matrix Multiplication Proof**

- (a) Given that matrix  $A$  is square and has linearly independent columns, which of the following are true?
- i.  $A$  is full rank
  - ii.  $A$  has a trivial nullspace
  - iii.  $A\vec{x} = \vec{b}$  has a unique solution for all  $\vec{b}$
  - iv.  $A$  is invertible
  - v. The determinant of  $A$  is non-zero

- (b) Let two square matrices  $M_1, M_2 \in \mathbb{R}^{2 \times 2}$  each have linearly independent columns. Prove that  $G = M_1 M_2$  also has linearly independent columns.

## 2. The Romulan Ruse

While scanning parts of the galaxy for alien civilization, the starship USS Enterprise NC-1701D encounters a Romulan starship that is known for advanced cloaking devices.

- (a) The Romulan illusion technology causes a point  $(x_0, y_0)$  to transform or *map* to  $(u_0, v_0)$ . Similarly,  $(x_1, y_1)$  is mapped to  $(u_1, v_1)$ . Figure 1 and Table 1 show these points.




Figure 1: Figure for part (a)

Original Point	Mapped Point
$(x_0, y_0) = (500, 500)$	$(u_0, v_0) = (500, 1500)$
Original Point	Mapped Point
$(x_1, y_1) = (1000, 500)$	$(u_1, v_1) = (1000, 1500)$

Table 1: Original and Mapped Points

**Find a transformation matrix  $\mathbf{A}_0$  such that**

$$\begin{bmatrix} u_0 \\ v_0 \end{bmatrix} = \mathbf{A}_0 \begin{bmatrix} x_0 \\ y_0 \end{bmatrix}, \text{ and } \begin{bmatrix} u_1 \\ v_1 \end{bmatrix} = \mathbf{A}_0 \begin{bmatrix} x_1 \\ y_1 \end{bmatrix}.$$

- (b) In this scenario, every point on the Romulan ship  $(x_m, y_m)$  is mapped to  $(u_m, v_m)$ , such that vector  $\begin{bmatrix} x_m \\ y_m \end{bmatrix}$  is rotated counterclockwise by  $30^\circ$  and then scaled by 2 in the x- and y-directions. This transformation is shown in Figure 2.




Figure 2: Figure for part (b)

$\theta$	$\sin \theta$	$\cos \theta$	$\tan \theta$
$0^\circ$	0	1	0
$30^\circ$	$\frac{1}{2}$	$\frac{\sqrt{3}}{2}$	$\frac{1}{\sqrt{3}}$
$45^\circ$	$\frac{1}{\sqrt{2}}$	$\frac{1}{\sqrt{2}}$	1
$60^\circ$	$\frac{\sqrt{3}}{2}$	$\frac{1}{2}$	$\sqrt{3}$
$90^\circ$	1	0	$\infty$

Table 2: Trigonometric Table

Find a transformation matrix  $\mathbf{R}$  such that  $\begin{bmatrix} u_m \\ v_m \end{bmatrix} = \mathbf{R} \begin{bmatrix} x_m \\ y_m \end{bmatrix}$ .

The Romulan ship has launched a probe into space and the Enterprise is trying to destroy the probe by firing a photon torpedo along a straight line from point  $(0,0)$  towards the probe.

- (c) The Romulan generals found a clever way to hide the probe by transforming (mapping) its position with a *cloaking* (transformation) matrix  $\mathbf{A}_p$ :

$$\mathbf{A}_p = \begin{bmatrix} 1 & 3 \\ 2 & 6 \end{bmatrix}.$$

They positioned the probe at  $(x_p, y_p)$  so that it maps to

$$(u_p, v_p) = (0, 0), \text{ where } \begin{bmatrix} u_p \\ v_p \end{bmatrix} = \mathbf{A}_p \begin{bmatrix} x_p \\ y_p \end{bmatrix}.$$

This scenario is shown in Figure 3. The initial position of the torpedo is  $(0,0)$  and the torpedo cannot be fired on its initial position! Impressive trick indeed!

**Find the possible positions of the probe  $(x_p, y_p)$  so that  $(u_p, v_p) = (0, 0)$ .**




Figure 3: Figure for part (c)

- (d) It turns out the Romulan engineers were not as smart as the Enterprise engineers. Their calculations did not work out and they positioned the probe at  $(x_q, y_q)$  such that the *cloaking* (transformation) matrix,  $\mathbf{A}_p$ , mapped it to  $(u_q, v_q)$ , where

$$\begin{bmatrix} u_q \\ v_q \end{bmatrix} = \mathbf{A}_p \begin{bmatrix} x_q \\ y_q \end{bmatrix}, \text{ and } \mathbf{A}_p = \begin{bmatrix} 1 & 3 \\ 2 & 6 \end{bmatrix}.$$

As a result, the torpedo, while traveling straight from  $(0, 0)$  to  $(u_q, v_q)$ , hit the probe at  $(x_q, y_q)$  on the way!

The scenario is shown in Figure 4. For the torpedo to

hit the probe, we must have  $\begin{bmatrix} u_q \\ v_q \end{bmatrix} = \lambda \begin{bmatrix} x_q \\ y_q \end{bmatrix}$ , where  $\lambda$

is a real number.

**Find the possible positions of the probe  $(x_q, y_q)$  so that  $(u_q, v_q) = (\lambda x_q, \lambda y_q)$ . Remember that the torpedo cannot be fired on  $(0, 0)$ . This means that  $(u_q, v_q) = (\lambda x_q, \lambda y_q)$  cannot be  $(0, 0)$ .**




Figure 4: Figure for part (d)