

Welcome to EECS 16A!

Designing Information Devices and Systems I

Ana Arias and Miki Lustig

Fall 2022

Lecture 5A
EigenVals/Vecs/Spaces



Announcements

- Last time:
 - Computing the determinant
 - Eigen Values and Eigen Vectors of a Matrix
 - Example via page-rank
- Today:
 - More on Eigenvalues, spaces and vectors

Recap

- What have we done in EECS 16A so far?
 1. Set of Equations
 2. Matrix vector multiplication
 3. Gaussian elimination
 4. Span, linear independence
 5. Matrices as transformations
 6. Matrix inversion
 7. Column space, null space
 8. Eigenvalues ; Eigenspace



Eigenvalues and Eigenvectors

$$A \vec{v} = \lambda \vec{v}$$

Must be square!

Any number!

Non-zero!

Q: What does $\lambda = 0$ mean?

$\vec{v} \neq \vec{0}$


Eigen Values and Eigen Vectors

- Definition: Let $A \in \mathbb{R}^{N \times N}$ be a square matrix, and $\lambda \in \mathbb{R}$ if $\exists \vec{v} \neq \vec{0}$ such that $A\vec{v} = \lambda\vec{v}$, then λ is an **eigenvalue** of A , \vec{v} is an **eigenvector** and $\text{Null}(A - \lambda I)$ is its **eigenspace**.

**In general $\lambda \in \mathbb{C}$

Disciplined Approach:

$$A \vec{v} = \lambda \vec{v}$$

1. Form $B_\lambda = A - \lambda I$
2. Find all the λ s resulting in a non-trivial null space for B_λ
 - Solve: $\det(B_\lambda) = 0$
 -  N^{th} order characteristic polynomial with N solutions
 - Each solution is an eigenvalue!
3. For each λ find the vector space $\text{Null}(B_\lambda)$

Solutions for the Characteristic Polynomial

$$A = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$$

$$\det(A - \lambda I) = \det \left(\begin{bmatrix} a - \lambda & b \\ c & d - \lambda \end{bmatrix} \right) = (a - \lambda)(d - \lambda) - bc = 0$$

$$\lambda^2 - (a + d)\lambda + (ad - bc) = 0$$

- Three cases:
 - Two real distinct eigenvalues
 - Single repeated eigenvalue
 - Two complex-valued eigenvalues

Distinct Eigenvalues

- Theorem: Let $A \in \mathbb{R}^{N \times N}$, with M distinct eigenvalues and corresponding eigenvectors $\lambda_i, \vec{v}_i \mid 1 \leq i \leq M$. It is the case that all \vec{v}_i are linearly independent. (Proof 9.6.2 in the notes)
- If $A \in \mathbb{R}^{2 \times 2}$ has two distinct eigenvalues, then:
 - \vec{v}_1, \vec{v}_2 are linearly independent
 - $\text{Span}\{\vec{v}_1, \vec{v}_2\} = \mathbb{R}^2$ — form a basis!

Proof 9.6.1 in the notes

Concept: By contradiction. Assume linear dependence \rightarrow This results in either $\lambda_1 = \lambda_2$, or $\vec{v}_2 = \vec{0}$

Matrix transformations


$$A \vec{v} = \lambda \vec{v}$$

What does the matrix do?

What is the A matrix?

What are its eigenvectors?

What are its eigenvalues?



Eigen Value Decomposition

$$A = \begin{bmatrix} 1 & 0 \\ 0 & 2 \end{bmatrix}$$

$$\det(A - \lambda I) = \begin{vmatrix} 1 - \lambda & 0 \\ 0 & 2 - \lambda \end{vmatrix} = (1 - \lambda)(2 - \lambda) - 0 = 0$$

$$\lambda_1 = 1$$

$$\lambda_2 = 2$$

$$\begin{bmatrix} 0 & 0 \\ 0 & 1 \end{bmatrix} \vec{v} = 0$$

$$\begin{bmatrix} -1 & 0 \\ 0 & 0 \end{bmatrix} \vec{v} = 0$$

$$\left[\begin{array}{cc|c} 0 & 1 & 0 \\ 0 & 0 & 0 \end{array} \right] \Rightarrow \begin{array}{l} v_2 = 0 \\ v_1 = \text{F.V.} \end{array}$$
$$\vec{v}_1 \in \text{Span}\left\{ \begin{bmatrix} 1 \\ 0 \end{bmatrix} \right\}$$

$$\left[\begin{array}{cc|c} 1 & 0 & 0 \\ 0 & 0 & 0 \end{array} \right] \Rightarrow \begin{array}{l} v_2 = \text{F.V.} \\ v_1 = 0 \end{array}$$
$$\vec{v}_2 \in \text{Span}\left\{ \begin{bmatrix} 0 \\ 1 \end{bmatrix} \right\}$$

Matrix transformations


$$A \vec{v} = \lambda \vec{v}$$

What does the matrix do?

What is the A matrix?

What are its eigenvectors?

What are its eigenvalues?



Eigenvectors as a basis

$$A = \begin{bmatrix} 1 & 0 \\ 0 & 2 \end{bmatrix} \quad \lambda_1 = 1, \lambda_2 = 2 \quad \vec{v}_1 = \begin{bmatrix} 1 \\ 0 \end{bmatrix} \quad \vec{v}_2 = \begin{bmatrix} 0 \\ 1 \end{bmatrix}$$

$$A\vec{v}_1 = 1 \cdot \vec{v}_1 \quad A\vec{v}_2 = 2 \cdot \vec{v}_2$$

Q: What about $\vec{v}_3 = \begin{bmatrix} 1 \\ 1 \end{bmatrix}$? $\vec{v}_3 = \begin{bmatrix} 1 \\ 1 \end{bmatrix} = 1 \cdot \vec{v}_1 + 1 \cdot \vec{v}_2$

$$\begin{aligned} A\vec{v}_3 &= A(1 \cdot \vec{v}_1 + 1 \cdot \vec{v}_2) = A\vec{v}_1 + A\vec{v}_2 \\ &= \vec{v}_1 + 2\vec{v}_2 \\ &= \begin{bmatrix} 1 \\ 0 \end{bmatrix} + 2 \begin{bmatrix} 0 \\ 1 \end{bmatrix} = \begin{bmatrix} 1 \\ 2 \end{bmatrix} \end{aligned}$$

Example from last time:

$$A = \begin{bmatrix} 1/2 & 0 \\ 1/2 & 1 \end{bmatrix} \quad \vec{v}_1 \in \text{Span} \left\{ \begin{bmatrix} 1 \\ -1 \end{bmatrix} \right\} \quad \vec{v}_2 \in \text{Span} \left\{ \begin{bmatrix} 0 \\ 1 \end{bmatrix} \right\}$$

$\lambda_1 = 1/2$ $\lambda_2 = 1$

- \vec{v}_1, \vec{v}_2 are linearly independent \rightarrow basis for \mathbb{R}^2

Q: What about $\vec{v}_3 = \begin{bmatrix} 2 \\ 2 \end{bmatrix}$? $\vec{v}_3 = \alpha \vec{v}_1 + \beta \vec{v}_2$



Example from last time:

$$A = \begin{bmatrix} 1/2 & 0 \\ 1/2 & 1 \end{bmatrix} \quad \vec{v}_1 \in \text{Span} \left\{ \begin{bmatrix} 1 \\ -1 \end{bmatrix} \right\} \quad \vec{v}_2 \in \text{Span} \left\{ \begin{bmatrix} 0 \\ 1 \end{bmatrix} \right\}$$

$\lambda_1 = 1/2$ $\lambda_2 = 1$

- \vec{v}_1, \vec{v}_2 are linearly independent \rightarrow basis for \mathbb{R}^2

Q: What about $\vec{v}_3 = \begin{bmatrix} 2 \\ 2 \end{bmatrix}$? $\vec{v}_3 = \alpha \vec{v}_1 + \beta \vec{v}_2$

$$\begin{bmatrix} 1 & 0 \\ -1 & 1 \end{bmatrix} \begin{bmatrix} \alpha \\ \beta \end{bmatrix} = \begin{bmatrix} 2 \\ 2 \end{bmatrix} \Rightarrow$$

Example from last time:

$$A = \begin{bmatrix} 1/2 & 0 \\ 1/2 & 1 \end{bmatrix} \quad \vec{v}_1 \in \text{Span} \left\{ \begin{bmatrix} 1 \\ -1 \end{bmatrix} \right\} \quad \vec{v}_2 \in \text{Span} \left\{ \begin{bmatrix} 0 \\ 1 \end{bmatrix} \right\}$$

$\lambda_1 = 1/2$ $\lambda_2 = 1$

- \vec{v}_1, \vec{v}_2 are linearly independent \rightarrow basis for \mathbb{R}^2

Q: What about $\vec{v}_3 = \begin{bmatrix} 2 \\ 2 \end{bmatrix}$? $\vec{v}_3 = \alpha \vec{v}_1 + \beta \vec{v}_2$

$$\begin{bmatrix} 1 & 0 \\ -1 & 1 \end{bmatrix} \begin{bmatrix} \alpha \\ \beta \end{bmatrix} = \begin{bmatrix} 2 \\ 2 \end{bmatrix} \Rightarrow \left[\begin{array}{cc|c} 1 & 0 & 2 \\ -1 & 1 & 2 \end{array} \right] \Rightarrow$$

Example from last time:

$$A = \begin{bmatrix} 1/2 & 0 \\ 1/2 & 1 \end{bmatrix} \quad \vec{v}_1 \in \text{Span} \left\{ \begin{bmatrix} 1 \\ -1 \end{bmatrix} \right\} \quad \vec{v}_2 \in \text{Span} \left\{ \begin{bmatrix} 0 \\ 1 \end{bmatrix} \right\}$$

$\lambda_1 = 1/2$ $\lambda_2 = 1$

- \vec{v}_1, \vec{v}_2 are linearly independent \rightarrow basis for \mathbb{R}^2

Q: What about $\vec{v}_3 = \begin{bmatrix} 2 \\ 2 \end{bmatrix}$? $\vec{v}_3 = \alpha \vec{v}_1 + \beta \vec{v}_2$

$$\begin{bmatrix} 1 & 0 \\ -1 & 1 \end{bmatrix} \begin{bmatrix} \alpha \\ \beta \end{bmatrix} = \begin{bmatrix} 2 \\ 2 \end{bmatrix} \Rightarrow \left[\begin{array}{cc|c} 1 & 0 & 2 \\ -1 & 1 & 2 \end{array} \right] \Rightarrow \left[\begin{array}{cc|c} 1 & 0 & 2 \\ 0 & 1 & 4 \end{array} \right] \Rightarrow$$

Example from last time:

$$A = \begin{bmatrix} 1/2 & 0 \\ 1/2 & 1 \end{bmatrix} \quad \vec{v}_1 \in \text{Span} \left\{ \begin{bmatrix} 1 \\ -1 \end{bmatrix} \right\} \quad \vec{v}_2 \in \text{Span} \left\{ \begin{bmatrix} 0 \\ 1 \end{bmatrix} \right\}$$

$\lambda_1 = 1/2$ $\lambda_2 = 1$

- \vec{v}_1, \vec{v}_2 are linearly independent \rightarrow basis for \mathbb{R}^2

Q: What about $\vec{v}_3 = \begin{bmatrix} 2 \\ 2 \end{bmatrix}$? $\vec{v}_3 = \alpha \vec{v}_1 + \beta \vec{v}_2$

$$\begin{bmatrix} 1 & 0 \\ -1 & 1 \end{bmatrix} \begin{bmatrix} \alpha \\ \beta \end{bmatrix} = \begin{bmatrix} 2 \\ 2 \end{bmatrix} \Rightarrow \begin{bmatrix} 1 & 0 & | & 2 \\ -1 & 1 & | & 2 \end{bmatrix} \Rightarrow \begin{bmatrix} 1 & 0 & | & 2 \\ 0 & 1 & | & 4 \end{bmatrix} \Rightarrow \begin{matrix} \alpha = 2 \\ \beta = 4 \end{matrix}$$

Example from last time:

$$A = \begin{bmatrix} 1/2 & 0 \\ 1/2 & 1 \end{bmatrix} \quad \vec{v}_1 \in \text{Span} \left\{ \begin{bmatrix} 1 \\ -1 \end{bmatrix} \right\} \quad \vec{v}_2 \in \text{Span} \left\{ \begin{bmatrix} 0 \\ 1 \end{bmatrix} \right\}$$

$\lambda_1 = 1/2$ $\lambda_2 = 1$

$$\begin{bmatrix} 1 & 0 \\ -1 & 1 \end{bmatrix} \begin{bmatrix} \alpha \\ \beta \end{bmatrix} = \begin{bmatrix} 2 \\ 2 \end{bmatrix} \Rightarrow \begin{bmatrix} 1 & 0 & | & 2 \\ -1 & 1 & | & 2 \end{bmatrix} \Rightarrow \begin{bmatrix} 1 & 0 & | & 2 \\ 0 & 1 & | & 4 \end{bmatrix} \Rightarrow \begin{matrix} \alpha = 2 \\ \beta = 4 \end{matrix}$$

$$\begin{aligned} A \vec{v}_3 &= A(2\vec{v}_1 + 4\vec{v}_2) = 2A\vec{v}_1 + 4A\vec{v}_2 \\ &= 2\left(\frac{1}{2}\vec{v}_1\right) + 4(1 \cdot \vec{v}_2) \\ &= \vec{v}_1 + 4\vec{v}_2 \\ &= \begin{bmatrix} 1 \\ -1 \end{bmatrix} + 4 \begin{bmatrix} 0 \\ 1 \end{bmatrix} = \begin{bmatrix} 1 \\ 3 \end{bmatrix} \end{aligned}$$

Matrix transformations


$$A \vec{v} = \lambda \vec{v}$$

What does the matrix do?

What is the A matrix?

What are its eigenvectors?

What are its eigenvalues?



Repeated EigenValues

$$A = \begin{bmatrix} 2 & 0 \\ 0 & 2 \end{bmatrix}$$

$$\det(A - \lambda I) = \begin{vmatrix} 2 - \lambda & 0 \\ 0 & 2 - \lambda \end{vmatrix} = (2 - \lambda)(2 - \lambda) - 0 = 0$$

$$\lambda_{1,2} = 2$$

$$\text{Null}(A - 2I) = \text{Null}(\vec{0}) = \mathbb{R}^2$$

Eigen space is 2 dimensional!

In general, multiplicity of Eigen-values will result in a multi-dimensional eigenspace

Except if the matrix is defective 😭

Repeated EigenValues


$$A \vec{v} = \lambda \vec{v}$$

What does the matrix do?

What is the A matrix?

What are its eigenvectors?

What are its eigenvalues?



Defective Matrices


$$A \vec{v} = \lambda \vec{v}$$

What does the matrix do?

What is the A matrix?

What are its eigenvectors?

What are its eigenvalues?



Defective Matrix

Outside of class scope 🤔

$$A = \begin{bmatrix} 1 & 1/4 \\ 0 & 1 \end{bmatrix}$$

$$\det(A - \lambda I) = \begin{vmatrix} 1 - \lambda & 1/4 \\ 0 & 1 - \lambda \end{vmatrix} = (1 - \lambda)(1 - \lambda) - 0 = 0$$

$$\lambda_{1,2} = 1$$

$$\text{Null}(A - I) = \text{Null} \left\{ \begin{bmatrix} 0 & 1/4 \\ 0 & 0 \end{bmatrix} \right\}$$

$$\vec{v}_1 \in \text{Span} \left\{ \begin{bmatrix} 1 \\ 0 \end{bmatrix} \right\}$$

Eigen space is only 1 dimensional!

Matrix is called defective 😭


Matrix transformations - Complex Eigenvalues

What does the matrix do?

What is the A matrix?

What are its eigenvectors?

What are its eigenvalues?




Matrix transformations

What does the matrix do?





What is the A matrix?


What are its eigenvectors?

What are its eigenvalues?



Back 2 PageRank

		From			
					
To		0	$\frac{1}{2}$	0	0
		$\frac{1}{3}$	0	0	$\frac{1}{2}$
		$\frac{1}{3}$	0	0	$\frac{1}{2}$
		$\frac{1}{3}$	$\frac{1}{2}$	1	0



Back 2 Page Rank


$$\vec{x}^* = \begin{bmatrix} 0.12 \\ 0.24 \\ 0.24 \\ 0.4 \end{bmatrix} = A \vec{x}^* = \begin{bmatrix} 0 & 1/2 & 0 & 0 \\ 1/3 & 0 & 0 & 1/2 \\ 1/3 & 0 & 0 & 1/2 \\ 1/3 & 1/2 & 1 & 0 \end{bmatrix} \begin{bmatrix} 0.12 \\ 0.24 \\ 0.24 \\ 0.4 \end{bmatrix}$$

steady state!



Back 2 PageRank

$$\vec{x}(t+1) = \begin{bmatrix} 0 & 1/2 & 0 & 0 \\ 1/3 & 0 & 0 & 1/2 \\ 1/3 & 0 & 0 & 1/2 \\ 1/3 & 1/2 & 1 & 0 \end{bmatrix} \vec{x}(t)$$



$\vec{x}(t) \Rightarrow$ Page ranking

$$\vec{x}(0) = \begin{bmatrix} 1/4 \\ 1/4 \\ 1/4 \\ 1/4 \end{bmatrix}$$

equal \Downarrow Ranking

$t=1$	$t=2$	$t=3$...	$t=100$
$\begin{bmatrix} 0.125 \\ 0.208 \\ 0.208 \\ 0.458 \end{bmatrix}$	$\begin{bmatrix} 0.104 \\ 0.271 \\ 0.271 \\ 0.354 \end{bmatrix}$	$\begin{bmatrix} 0.135 \\ 0.212 \\ 0.212 \\ 0.441 \end{bmatrix}$...	$\begin{bmatrix} 0.12 \\ 0.24 \\ 0.24 \\ 0.4 \end{bmatrix}$



General Initialization for a Transition Matrix System

$$\vec{x}(t + 1) = A \vec{x}(t)$$

Assume $\lambda_i \mid 1 \leq i \leq N$ are distinct $\Rightarrow \text{Span}\{\vec{v}_i \mid 1 \leq i \leq N\} = \mathbb{R}^N$

$$\begin{aligned}\vec{x}(1) &= A \vec{x}(0) \\ &= A(\alpha_1 \vec{v}_1 + \alpha_2 \vec{v}_2 + \cdots + \alpha_N \vec{v}_N) \\ &= \alpha_1 A \vec{v}_1 + \alpha_2 A \vec{v}_2 + \cdots + \alpha_N A \vec{v}_N \\ &= \alpha_1 \lambda_1 \vec{v}_1 + \alpha_2 \lambda_2 \vec{v}_2 + \cdots + \alpha_N \lambda_N \vec{v}_N\end{aligned}$$

$$\begin{aligned}\vec{x}(2) &= A \vec{x}(1) \\ &= A(\alpha_1 \lambda_1 \vec{v}_1 + \alpha_2 \lambda_2 \vec{v}_2 + \cdots + \alpha_N \lambda_N \vec{v}_N) \\ &= \alpha_1 \lambda_1^2 \vec{v}_1 + \alpha_2 \lambda_2^2 \vec{v}_2 + \cdots + \alpha_N \lambda_N^2 \vec{v}_N\end{aligned}$$

General Initialization for a Transition Matrix System

$$\vec{x}(t+1) = A\vec{x}(t)$$

Assume $\lambda_i \mid 1 \leq i \leq N$ are distinct $\Rightarrow \text{Span}\{\vec{v}_i \mid 1 \leq i \leq N\} = \mathbb{R}^N$

$$\vec{x}(2) = A\vec{x}(1)$$

$$= A(\alpha_1\lambda_1\vec{v}_1 + \alpha_2\lambda_2\vec{v}_2 + \cdots + \alpha_N\lambda_N\vec{v}_N)$$

$$= \alpha_1\lambda_1^2\vec{v}_1 + \alpha_2\lambda_2^2\vec{v}_2 + \cdots + \alpha_N\lambda_N^2\vec{v}_N$$

•
•
•

$$\vec{x}(t) = \alpha_1\lambda_1^t\vec{v}_1 + \alpha_2\lambda_2^t\vec{v}_2 + \cdots + \alpha_N\lambda_N^t\vec{v}_N$$

$$\lim_{t \rightarrow \infty} \vec{x}(t) = ?$$

$$\lim_{t \rightarrow \infty} \lambda_i^t = \begin{cases} 0 & , \quad |\lambda| < 1 \\ 1 & , \quad \lambda = 1 \\ (-1)^t & , \quad \lambda = -1 \\ \infty & , \quad |\lambda| > 1 \end{cases}$$

Back 2 PageRank

$$A = \begin{bmatrix} 0 & 1/2 & 0 & 0 \\ 1/3 & 0 & 0 & 1/2 \\ 1/3 & 0 & 0 & 1/2 \\ 1/3 & 1/2 & 1 & 0 \end{bmatrix}$$

$$\lambda_1 = 1$$

$$\lambda_2 = -0.092$$

$$\lambda_3 = -0.91$$

$$\lambda_4 = 0$$

$$\vec{v}_1 = \begin{bmatrix} 0.12 \\ 0.24 \\ 0.24 \\ 0.4 \end{bmatrix}$$

$$\vec{v}_2 = \begin{bmatrix} 0.44 \\ -0.08 \\ -0.08 \\ -0.28 \end{bmatrix}$$

$$\vec{v}_3 = \begin{bmatrix} -0.14 \\ 0.26 \\ 0.26 \\ -0.37 \end{bmatrix}$$

$$\vec{v}_4 = \begin{bmatrix} 0.43 \\ 0 \\ -0.14 \\ -0.29 \end{bmatrix}$$

$$\vec{x}(t) = A^t \vec{x}(0)$$

Back 2 PageRank

$\lambda_1 = 1$

$\lambda_2 = -0.092$

$\lambda_3 = -0.091$

$\lambda_4 = 0$

$$\vec{v}_1 = \begin{bmatrix} 0.12 \\ 0.24 \\ 0.24 \\ 0.4 \end{bmatrix}$$

$$\vec{v}_2 = \begin{bmatrix} 0.44 \\ -0.08 \\ -0.08 \\ -0.28 \end{bmatrix}$$

$$\vec{v}_3 = \begin{bmatrix} -0.14 \\ 0.26 \\ 0.26 \\ -0.37 \end{bmatrix}$$

$$\vec{v}_4 = \begin{bmatrix} 0.43 \\ 0 \\ -0.14 \\ -0.29 \end{bmatrix}$$

$$\vec{x}_0 = \begin{bmatrix} 0.25 \\ 0.25 \\ 0.25 \\ 0.25 \end{bmatrix} = \alpha_1 \vec{v}_1 + \alpha_2 \vec{v}_2 + \alpha_3 \vec{v}_3 + \alpha_4 \vec{v}_4$$

$$\begin{bmatrix} \vec{v}_1 & \vec{v}_2 & \vec{v}_3 & \vec{v}_4 \end{bmatrix} \begin{bmatrix} \alpha_1 \\ \alpha_2 \\ \alpha_3 \\ \alpha_4 \end{bmatrix} = \begin{bmatrix} 0.25 \\ 0.25 \\ 0.25 \\ 0.25 \end{bmatrix}$$

V $\vec{\alpha}$ x_0

$$V \vec{\alpha} = \vec{x}_0$$

(1) Solve via G.E

$$\vec{\alpha} = V^{-1} \vec{x}_0$$

$$\vec{\alpha} = \begin{bmatrix} 1 \\ 0.34 \\ 0.15 \\ 0 \end{bmatrix}$$


$$\vec{x}(t) = A^t \vec{x}(0)$$

$$\begin{aligned} A^t \vec{x}(0) &= A(1 \vec{v}_1 + 0.34 \vec{v}_2 + 0.15 \vec{v}_3 + 0 \vec{v}_4) \\ &= 1 \cdot 1^t \vec{v}_1 + 0.34(-0.092)^t \vec{v}_2 + 0.15(-0.91)^t \vec{v}_3 + 0 \cdot 0^t \vec{v}_4 \end{aligned}$$

$$\lim_{t \rightarrow \infty} A^t \vec{x}(0) = \vec{v}_1$$

Back to Lab — Single Pixel Camera

- What are the best patterns?




Imaging Model and Reconstruction

Measurement mask/matrix

$$\vec{y} = A \vec{x}$$

Sensor data

Unknown image




We saw that it is possible to come up with a system that has A^{-1}

So,
$$\vec{x} = A^{-1} \vec{y}$$

Non-ideal imaging

Measurement mask/matrix



We saw that it is possible to come up with a system that has A^{-1}

So,
$$\vec{x} = A^{-1}\vec{y} - A^{-1}\vec{w}$$

Reconstruction error

$$A^{-1}\vec{w} = \alpha_1\lambda_1\vec{v}_1 + \alpha_2\lambda_2\vec{v}_2 + \dots + \alpha_N\lambda_N\vec{v}_N$$

Want to design A , such that A^{-1} has small eigenvalues!

Design of a Reflection matrix


Design a reflection matrix around the vector $\begin{bmatrix} 2 \\ 1 \end{bmatrix}$?

Q: What are the eigenvectors?

A: $\vec{v}_1 = \begin{bmatrix} 2 \\ 1 \end{bmatrix}, \vec{v}_2 = \begin{bmatrix} -1 \\ 2 \end{bmatrix}$

Q: What are the eigenvalues?

A: $\lambda_1 = 1, \lambda_2 = -1$



Designing a matrix with specific Eigenvals/vecs

We know:

$$A \vec{v} = \lambda \vec{v}$$

$$\vec{v}_1 = \begin{bmatrix} 2 \\ 1 \end{bmatrix}, \vec{v}_2 = \begin{bmatrix} -1 \\ 2 \end{bmatrix}$$

$$\lambda_1 = 1, \lambda_2 = -1$$

Set linear equations:

$$\begin{bmatrix} a_1 & a_2 \\ a_3 & a_4 \end{bmatrix} \begin{bmatrix} 2 \\ 1 \end{bmatrix} = \begin{bmatrix} 2 \\ 1 \end{bmatrix}$$

$$\begin{bmatrix} a_1 & a_2 \\ a_3 & a_4 \end{bmatrix} \begin{bmatrix} -1 \\ 2 \end{bmatrix} = \begin{bmatrix} -1 \\ -2 \end{bmatrix}$$

unknown

$$\begin{bmatrix} \\ \\ \\ \end{bmatrix} \begin{bmatrix} a_1 \\ a_2 \\ a_3 \\ a_4 \end{bmatrix} = \begin{bmatrix} \\ \\ \\ \end{bmatrix}$$

Designing a matrix with specific Eigenvals/vecs

We know:

$$A \vec{v} = \lambda \vec{v}$$

$$\vec{v}_1 = \begin{bmatrix} 2 \\ 1 \end{bmatrix}, \vec{v}_2 = \begin{bmatrix} -1 \\ 2 \end{bmatrix}$$

$$\lambda_1 = 1, \lambda_2 = -1$$

Set linear equations:

$$\begin{bmatrix} a_1 & a_2 \\ a_3 & a_4 \end{bmatrix} \begin{bmatrix} 2 \\ 1 \end{bmatrix} = \begin{bmatrix} 2 \\ 1 \end{bmatrix}$$

$$\begin{bmatrix} a_1 & a_2 \\ a_3 & a_4 \end{bmatrix} \begin{bmatrix} -1 \\ 2 \end{bmatrix} = \begin{bmatrix} -1 \\ -2 \end{bmatrix}$$

unknown

$$\begin{bmatrix} 2 & 1 & 0 & 0 \\ 0 & 0 & 2 & 1 \\ -1 & 2 & 0 & 0 \\ 0 & 0 & -1 & 2 \end{bmatrix} \begin{bmatrix} a_1 \\ a_2 \\ a_3 \\ a_4 \end{bmatrix} = \begin{bmatrix} 2 \\ 1 \\ 1 \\ -2 \end{bmatrix}$$

G.E. \Rightarrow

$$A = \begin{bmatrix} 0.6 & 0.8 \\ 0.8 & -0.6 \end{bmatrix}$$