
EECS 16A Designing Information Devices and Systems I
Spring 2022 Discussion 12A

1. Inner Product Properties

For this question, we will verify our coordinate definition of the inner product

$$\langle \vec{x}, \vec{y} \rangle = x_1y_1 + x_2y_2 + \dots + x_ny_n, \quad \text{for any } \vec{x}, \vec{y} \in \mathbb{R}^n$$

indeed satisfies the key properties required for all inner products, but presently for the 2-dimensional case. Suppose $\vec{x}, \vec{y}, \vec{z} \in \mathbb{R}^2$ for the following parts:

- Show symmetry $\langle \vec{x}, \vec{y} \rangle = \langle \vec{y}, \vec{x} \rangle$.
- Show linearity $\langle \vec{x}, c\vec{y} + d\vec{z} \rangle = c\langle \vec{x}, \vec{y} \rangle + d\langle \vec{x}, \vec{z} \rangle$, where $c, d \in \mathbb{R}$ are real numbers.
- Show non-negativity $\langle \vec{x}, \vec{x} \rangle \geq 0$, with equality if and only if $\vec{x} = \vec{0}$.


2. Geometric Interpretation of the Inner Product

In this problem, we explore the geometric interpretation of the Euclidean inner product, restricting ourselves to vectors in \mathbb{R}^2 .

Remember that the formula for the inner product of two vectors can be expressed in terms of their magnitudes and the angle between them as follows:

$$\langle \vec{x}, \vec{y} \rangle = \|\vec{x}\| \|\vec{y}\| \cdot \cos \theta$$

The figure below may be helpful in illustrating this property:




For each sub-part, give an example of any two (nonzero) vectors $\vec{x}, \vec{y} \in \mathbb{R}^2$ that satisfy the stated condition and compute their inner product.

- (a) Give an example of a pair of parallel vectors (vectors that point in the same direction and have an angle of 0 degrees between them).
- (b) Give an example of a pair of anti-parallel vectors (vectors that point in opposite directions).
- (c) Give an example of a pair of perpendicular vectors (vectors that have an angle of 90 degrees between them).


3. Correlation

- (a) You are given the following two signals:



Sketch the linear cross-correlation of signal 1 with signal 2, that is find: $\text{corr}(\vec{s}_1, \vec{s}_2)[n]$ for $n = 0, 1, \dots, 4$. Do not assume the signals are periodic.

- (b) Now the pattern in \vec{s}_1 is repeated three times:



Sketch the linear cross-correlation of signal 1 with signal 2, $\text{corr}(\vec{s}_1, \vec{s}_2)[n]$, for $n = 0, 1, \dots, 4$.