

# Discussion 14A

The following notes are useful for this discussion: [Note 18](#).

## 1. Jacobians and Linear Approximation

Recall that for a scalar-valued function  $f(\vec{x}, \vec{y}) : \mathbb{R}^n \times \mathbb{R}^k \rightarrow \mathbb{R}$  with vector-valued arguments, we can linearize the function at  $(\vec{x}_*, \vec{y}_*)$ :

$$\hat{f}(\vec{x}, \vec{y}) = f(\vec{x}_*, \vec{y}_*) + \sum_{i=1}^n \frac{\partial f(\vec{x}_*, \vec{y}_*)}{\partial x_i} (x_i - x_{i,*}) + \sum_{j=1}^k \frac{\partial f(\vec{x}_*, \vec{y}_*)}{\partial y_j} (y_j - y_{j,*}). \quad (1)$$

In order to simplify this equation, we can define the following two vector quantities:

$$J_{\vec{x}}f = \begin{bmatrix} \frac{\partial f}{\partial x_1} & \cdots & \frac{\partial f}{\partial x_n} \end{bmatrix} \quad (2)$$

$$J_{\vec{y}}f = \begin{bmatrix} \frac{\partial f}{\partial y_1} & \cdots & \frac{\partial f}{\partial y_k} \end{bmatrix} \quad (3)$$

- (a) When the function  $\vec{f}(\vec{x}, \vec{y}) : \mathbb{R}^n \times \mathbb{R}^k \rightarrow \mathbb{R}^m$  takes in vectors and outputs a *vector* (rather than a scalar), we can view each dimension in  $\vec{f}$  independently as a separate function  $f_i$ , and linearize each of them as above:

$$\hat{\vec{f}}(\vec{x}, \vec{y}) = \begin{bmatrix} \hat{f}_1(\vec{x}, \vec{y}) \\ \hat{f}_2(\vec{x}, \vec{y}) \\ \vdots \\ \hat{f}_m(\vec{x}, \vec{y}) \end{bmatrix} = \begin{bmatrix} f_1(\vec{x}_*, \vec{y}_*) + J_{\vec{x}}f_1 \cdot (\vec{x} - \vec{x}_*) + J_{\vec{y}}f_1 \cdot (\vec{y} - \vec{y}_*) \\ f_2(\vec{x}_*, \vec{y}_*) + J_{\vec{x}}f_2 \cdot (\vec{x} - \vec{x}_*) + J_{\vec{y}}f_2 \cdot (\vec{y} - \vec{y}_*) \\ \vdots \\ f_m(\vec{x}_*, \vec{y}_*) + J_{\vec{x}}f_m \cdot (\vec{x} - \vec{x}_*) + J_{\vec{y}}f_m \cdot (\vec{y} - \vec{y}_*) \end{bmatrix} \quad (4)$$

We can rewrite this in a clean way with the *Jacobian* of a vector-valued function:

$$J_{\vec{x}}\vec{f} = \begin{bmatrix} J_{\vec{x}}f_1 \\ J_{\vec{x}}f_2 \\ \vdots \\ J_{\vec{x}}f_m \end{bmatrix} = \begin{bmatrix} \frac{\partial f_1}{\partial x_1} & \cdots & \frac{\partial f_1}{\partial x_n} \\ \vdots & \ddots & \vdots \\ \frac{\partial f_m}{\partial x_1} & \cdots & \frac{\partial f_m}{\partial x_n} \end{bmatrix}, \quad (5)$$

and similarly

$$J_{\vec{y}}\vec{f} = \begin{bmatrix} \frac{\partial f_1}{\partial y_1} & \cdots & \frac{\partial f_1}{\partial y_k} \\ \vdots & \ddots & \vdots \\ \frac{\partial f_m}{\partial y_1} & \cdots & \frac{\partial f_m}{\partial y_k} \end{bmatrix}. \quad (6)$$

Then, the linearization becomes

$$\hat{\vec{f}}(\vec{x}, \vec{y}) = \vec{f}(\vec{x}_*, \vec{y}_*) + J_{\vec{x}}\vec{f}(\vec{x}_*, \vec{y}_*) \cdot (\vec{x} - \vec{x}_*) + J_{\vec{y}}\vec{f}(\vec{x}_*, \vec{y}_*) \cdot (\vec{y} - \vec{y}_*). \quad (7)$$

Let  $\vec{x} = \begin{bmatrix} x_1 \\ x_2 \end{bmatrix}$  and  $\vec{f}(\vec{x}) = \begin{bmatrix} x_1^2 x_2 \\ x_1 x_2^2 \end{bmatrix}$ . Find  $J_{\vec{x}}\vec{f}$ , applying the definition above.

(b) Evaluate the approximation of  $\vec{f}$  using  $\vec{x}_* = \begin{bmatrix} 2 \\ 3 \end{bmatrix}$  at the point  $\begin{bmatrix} 2.01 \\ 3.01 \end{bmatrix}$ , and compare with  $\vec{f}\left(\begin{bmatrix} 2.01 \\ 3.01 \end{bmatrix}\right)$ . Recall the definition that  $\vec{f}(\vec{x}) = \begin{bmatrix} x_1^2 x_2 \\ x_1 x_2^2 \end{bmatrix}$ .

(c) Let  $\vec{x}$  and  $\vec{y}$  be vectors with 2 rows, and let  $\vec{w}$  be another vector with 2 rows. Let  $\vec{f}(\vec{x}, \vec{y}) = \vec{x}\vec{y}^\top \vec{w}$ . Find  $J_{\vec{x}}\vec{f}$  and  $J_{\vec{y}}\vec{f}$ .


(d) (PRACTICE) Continuing the above part, find the linear approximation of  $\vec{f}$  near  $\vec{x} = \vec{y} = \begin{bmatrix} 1 \\ 1 \end{bmatrix}$  and with  $\vec{w} = \begin{bmatrix} 2 \\ 1 \end{bmatrix}$ .

## 2. Linearizing a Two-state System

We have a two-state nonlinear system defined by the following differential equation:

$$\frac{d}{dt} \begin{bmatrix} \beta(t) \\ \gamma(t) \end{bmatrix} = \frac{d}{dt} \vec{x}(t) = \begin{bmatrix} -2\beta(t) + \gamma(t) \\ g(\gamma(t)) + u(t) \end{bmatrix} = \vec{f}(\vec{x}(t), u(t)) \quad (8)$$

where  $\vec{x}(t) = \begin{bmatrix} \beta(t) \\ \gamma(t) \end{bmatrix}$  and  $g(\cdot)$  is a nonlinear function with the following graph:



The  $g(\cdot)$  is the only nonlinearity in this system. We want to linearize this entire system around a operating point/equilibrium. Any point  $\vec{x}_*$  is an operating point if  $\vec{f}(\vec{x}_*(t), u_*(t)) = \vec{0}$ .

(a) If we have fixed  $u_*(t) = -1$ , what values of  $\gamma$  and  $\beta$  will ensure  $\frac{d}{dt} \vec{x}(t) = \vec{f}(\vec{x}(t), u(t)) = \vec{0}$ ?



(b) Now that you have the three operating points, **linearize the system about the operating point**  $(\vec{x}_3^*, u_*)$  (that which has the largest value for  $\gamma$ ). Specifically, what we want is as follows. Let  $\delta \vec{x}_i(t) = \vec{x}(t) - \vec{x}_i^*$  for  $i = 1, 2, 3$ , and  $\delta u(t) = u(t) - u_*$ . We can in principle write the linearized system for each operating point in the following form:

$$\text{(linearization about } (\vec{x}_i^*, u_*)) \quad \frac{d}{dt} \delta \vec{x}_i(t) = A_i \delta \vec{x}_i(t) + B_i \delta u(t) + \vec{w}_i(t) \quad (9)$$

where  $\vec{w}_i(t)$  is a disturbance that also includes the approximation error due to linearization.

For this part, **find**  $A_3$  **and**  $B_3$ .

We have provided below the function  $g(\gamma)$  and its derivative  $\frac{\partial g}{\partial \gamma}$ .



(c) Which of the operating points are stable? Which are unstable?

**Contributors:**

- Neelesh Ramachandran.
- Kuan-Yun Lee.
- Alex Devonport.
- Kumar Krishna Agrawal.